

Allenamento contro resistenza per anziani: Position Statement della National Strength and Conditioning Association

Bibliografia

1. American College of Sports Medicine. Osteoporosis and exercise. *Med Sci Sports Exerc* 27: i-vii, 1995.
2. ACSM's Guidelines for Exercise Testing and Prescription. Philadelphia, PA: Lippincott William & Wilkins, 2009. pp. 400.
3. Aagaard P, Magnusson PS, Larsson B, Kjaer M, Krstrup P. Mechanical muscle function, morphology, and fiber type in lifelong trained elderly. *Med Sci Sports Exerc* 39: 1989-1996, 2007.
4. Aagaard P, Suetta C, Caserotti P, Magnusson SP, Kjaer M. Role of the nervous system in sarcopenia and muscle atrophy with aging: Strength training as a countermeasure. *Scand J Med Sci Sports* 20: 49-64, 2010.
5. Abney M, McPeck MS, Ober C. Broad and narrow heritabilities of quantitative traits in a founder population. *Am J Hum Genet* 68: 1302-1307, 2001.
6. Abraham O, Rodrigues RP, Nascimento VC, et al. Single- and multiple-set resistance training improves skeletal and respiratory muscle strength in elderly women. *Clin Interv Aging* 9: 1775-1782, 2014.
7. Ades PA, Ballor DL, Ashikaga T, Utton JL, Nair KS. Weight training improves walking endurance in healthy elderly persons. *Ann Intern Med* 124: 568-572, 1996.
8. International Diabetes Federation. Global Picture. In: 8th, ed. *IDF Diabetes Atlas*. Brussels, Belgium: International Diabetes Federation, 2017. pp. 40-65.
9. Ahlqvist A, Nyfors H, Suhonen R. Factors associated with older people's independent living from the viewpoint of health and functional capacity: A register-based study. *Nurs Open* 3: 79-89, 2016.
10. Ahtiainen JP, Hulmi JJ, Kraemer WJ, et al. Heavy resistance exercise training and skeletal muscle androgen receptor expression in younger and older men. *Steroids* 76: 183-192, 2011.
11. Ahtiainen JP, Hulmi JJ, Lehti M, et al. Effects of resistance training on expression of IGF-1 splice variants in younger and older men. *Eur J Sport Sci* 16: 1055-1063, 2016.
12. Ahtiainen JP, Nyman K, Huhtaniemi I, et al. Effects of resistance training on testosterone metabolism in younger and older men. *Exp Gerontol* 69: 148-158, 2015.
13. Alfaro-AcoHaA, Al Snih S, Raji MA, Kuo YF, Markides KS, Ottenbacher KJ. Handgrip strength and cognitive decline in older Mexican Americans. *J Gerontol A Biol Sci Med Sci* 61: 859-865, 2006.
14. Alley DE, Shardell MD, Peters KW, et al. Grip strength cutpoints for the identification of clinically relevant weakness. *J Gerontol A Biol Sci Med Sci* 69: 559-566, 2014.
15. Alvarez KJ, Kirchner S, Chu S, et al. Falls reduction and exercise training in an assisted living population. *J Aging Res* 2015: 957598, 2015.
16. American College of Sports Medicine. American College of Sports Medicine position stand: Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 41: 687-708, 2009.
17. American College of Sports Medicine. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 41: 687-708, 2009.
18. Balady GJ, Foster C, Chaitman B, Froelicher ES. Position stand: automated external defibrillators in health/fitness facilities. *Med Sci Sports Exerc* 34: 561-564, 2002.
19. American Geriatrics Society Panel on Exercise and Osteoarthritis. Exercise prescription for older adults with osteoarthritis pain: Consensus practice recommendations. A supplement to the AGS Clinical Practice Guidelines on the management of chronic pain in older adults. *J Am Geriatr Soc* 49: 808-823, 2001.
20. Amiya E, Taya M. Is exercise training appropriate for patients with advanced heart failure receiving continuous inotropic infusion? A review. *Clin Med Insights Cardiol* 12: 1179546817751438, 2018.
21. Amonette WE, English KL, Kraemer WJ. Evidence-Based Practice In Exercise Science: The Six-Step Approach. Champaign, IL: Human Kinetics, 2016.
22. Anderson DE, D'Agostino JM, Bruno AG, et al. Variations of CT-based trunk muscle attenuation by age, sex, and specific muscle. *J Gerontol A Biol Sci Med Sci* 68: 317-323, 2013.
23. Anthony K, Robinson K, Logan P, et al. Chair-based exercises for frail older people: A systematic review. *Biomed Res Int* 2013: 309506, 2013.
24. Arnold CM, Sran MM, Harrison EL. Exercise for fall risk reduction in community-dwelling older adults: A systematic review. *Physiother Can* 60: 358-372, 2008.
25. Artero EG, Ruiz JR, Ortega FB, et al. Muscular and cardiorespiratory fitness are independently associated with metabolic risk in adolescents: The HELENA study. *Pediatr Diabetes* 12: 704-712, 2011.
26. Balagopal P, Rooyackers OE, Adey DB, Ades PA, Nair KS. Effects of aging on in vivo synthesis of skeletal muscle myosin heavy-chain and sarco-plasmic protein in humans. *Am J Physiol* 273(4 Pt 1): E790-E800, 1997.
27. Balboa-Castillo T, Guallar-Castillon P, Leon-Munoz LM, et al. Physical activity and mortality related to obesity and functional status in older adults in Spain. *Am J Prev Med* 40: 39-46, 2011.
28. Ballor DL, Harvey-Berino JR, Ades PA, Cryan J, Calles-Escandon J. Contrasting effects of resistance and aerobic training on body composition and metabolism after diet-induced weight loss. *Metabolism* 45: 179-183, 1996.
29. Bamman MM, Hill VJ, Adams GR, et al. Gender differences in resistance-training-induced myofiber hypertrophy among older adults. *J Gerontol A Biol Sci Med Sci* 58: 108-116, 2003.
30. Banteen-Roche K, Xue QL, Ferrucci L, et al. Phenotype of frailty: Characterization in the women's health and aging studies. *J Gerontol A Biol Sci Med Sci* 61: 262-266, 2006.
31. Barber LA, Read F, Lovatt Stern J, Lichtwark G, Boyd RN. Medial gastrocnemius muscle volume in ambulant children with unilateral and bilateral cerebral palsy aged 2 to 9 years. *Dev Med Child Neurol* 58: 1146-1152, 2016.
32. Bardenheier B, Lin J, Zhuo X, et al. Disability-free life-years lost among adults aged 50 Years, with and without diabetes. *Diabetes Care* 39: 1222-1229, 2016.
33. Barnett A, Smith B, Lord SR, Williams M, Baumand A. Community-based group exercise improves balance and reduces falls in at-risk older people: A randomised controlled trial. *Age Ageing* 32: 407-414, 2003.
34. Barry BK, Warman GE, Carson RG. Age-related differences in rapid muscle activation after rate of force development training of the elbow flexors. *Exp Brain Res* 162: 122-132, 2005.

35. Barthalos I, Dorgo S, Kopka N, Plachy J, et al. Randomized controlled resistance training based physical activity trial for central European nursing home residing older adults. *J Sports Med Phys Fitness* 56: 1249–1257, 2016.
36. Basaria S. Reproductive aging in men. *Endocrinol Metab Clin North Am* 42: 255–270, 2013.
37. Bassey EJ, Fiatarone MA, O'Neill EF, et al. Leg extensor power and functional performance in very old men and women. *Clin Sci (Lond)* 82: 321–327, 1992.
38. Baum EE, Jarjoura D, Polen AE, Faur D, Rutecki G. Effectiveness of a group exercise program in a long-term care facility: A randomized pilot trial. *J Am Med Dir Assoc* 4: 74–80, 2003.
39. Baumgartner RN, Koehler KM, Gallagher D, et al. Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol* 147: 755–763, 1998.
40. Bautmans I, Njemini R, Vasseur S, et al. Biochemical changes in response to intensive resistance exercise training in the elderly. *Gerontology* 51: 253–265, 2005.
41. Baylis D, Ntani G, Edwards MH, et al. Inflammation, telomere length, and grip strength: A 10-year longitudinal study. *Calcif Tissue Int* 95: 54–63, 2014.
42. Bean JF, Herman S, Kiely DK, et al. Increased Velocity Exercise Specific to Task (INVEST) training: A pilot study exploring effects on leg power, balance, and mobility in community-dwelling older women. *J Am Geriatr Soc* 52: 799–804, 2004.
43. Bean JF, Kiely DK, Herman S, et al. The relationship between leg power and physical performance in mobility-limited older people. *J Am Geriatr Soc* 50: 461–467, 2002.
44. Bean JF, Kiely DK, LaRose S, et al. Increased velocity exercise specific to task training versus the National Institute on Aging's strength training program: Changes in limb power and mobility. *J Gerontol A Biol Sci Med Sci* 64: 983–991, 2009.
45. Bean JF, Leveille SG, Kiely DK, et al. A comparison of leg power and leg strength within the InCHIANTI study: Which influences mobility more? *J Gerontol A Biol Sci Med Sci* 58: 728–733, 2003.
46. Beasley LE, Koster A, Newman AB, et al. Inflammation and race and gender differences in computerized tomography-measured adipose depots. *Obesity (Silver Spring)* 17: 1062–1069, 2009.
47. Beck BR, Daly RM, Singh MA, Taaffe DR. Exercise and Sports Science Australia (ESSA) position statement on exercise prescription for the prevention and management of osteoporosis. *J Sci Med Sport* 20: 438–445, 2017.
48. Beck LH. The aging kidney. Defending a delicate balance of fluid and electrolytes. *Geriatrics* 55: 26–28, 2000; 31–2.
49. Beltran-Sanchez H, Jimenez MP, Subramanian SV. Assessing morbidity compression in two cohorts from the Health and Retirement Study. *J Epidemiol Community Health* 70: 1011–1016, 2016.
50. Bemben DA, Fettes NL, Bemben MG, Nabavi N, Koh ET. Musculo-skeletal responses to high- and low-intensity resistance training in early postmenopausal women. *Med Sci Sports Exerc* 32: 1949–1957, 2000.
51. Bemben MG, Massey BH, Bemben DA, Misner JE, Boileau RA. Isometric muscle force production as a function of age in healthy 20- to 74-year-old men. *Med Sci Sports Exerc* 23: 1302–1310, 1991.
52. Ben-Shlomo Y, Cooper R, Kuh D. The last two decades of life course epidemiology, and its relevance for research on ageing. *Int J Epidemiol* 45: 973–988, 2016.
53. Beneka A, Malliou P, Fatouros I, et al. Resistance training effects on muscular strength of elderly are related to intensity and gender. *J Sci Med Sport* 8: 274–283, 2005.
54. Bento PC, Pereira G, Ugrinowitsch C, Rodacki AL. Peak torque and rate of torque development in elderly with and without fall history. *Clin Biomech (Bristol, Avon)* 25: 450–454, 2010.
55. Bergman H, Ferrucci L, Guralnik J, et al. Frailty: An emerging research and clinical paradigm—issues and controversies. *J Gerontol A Biol Sci Med Sci* 62: 731–737, 2007.
56. Berzlanovich AM, Schopfer J, Keil W. Deaths due to physical restraint. *Dtsch Arztebl Int* 109: 27–32, 2012.
57. Bessa B, de Oliveira Leal V, Moraes C, et al. Resistance training in hemodialysis patients: A review. *Rehabil Nurs* 40: 111–126, 2015.
58. Beyer N, Suetta C. Older patients should be offered strength training early post surgery [in Danish]. *Ugeskr Laeger* 175: 2421–2424, 2013.
59. Bhattacharya RK, Bhattacharya SB. Late-onset hypogonadism and testosterone replacement in older men. *Clin Geriatr Med* 31: 631–644, 2015.
60. Bilodeau M, Henderson TK, Nolte BE, Pursley PJ, Sandfort GL. Effect of aging on fatigue characteristics of elbow flexor muscles during sustained submaximal contraction. *J Appl Physiol* (1985) 91: 2654–2664, 2001.
61. Binder EF, Birge SJ, Spina R, et al. Peak aerobic power is an important component of physical performance in older women. *J Gerontol A Biol Sci Med Sci* 54: M353–M356, 1999.
62. Binder EF, Schechtman KB, Ehsani AA, et al. Effects of exercise training on frailty in community-dwelling older adults: Results of a randomized, controlled trial. *J Am Geriatr Soc* 50: 1921–1928, 2002.
63. Binder EF, Yarasheski KE, Steger-May K, et al. Effects of progressive resistance training on body composition in frail older adults: Results of a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci* 60: 1425–1431, 2005.
64. Blazer DG. Depression in late life: Review and commentary. *J Gerontol A Biol Sci Med Sci* 58: 249–265, 2003.
65. Bodart G, Goffinet L, Morrhaye G, et al. Somatotrope GHRH/GH/IGF-1 axis at the crossroads between immunosenescence and frailty. *Ann N Y Acad Sci* 1351: 61–67, 2015.
66. Borde R, Hortobağyi T, Granacher U. Dose-response relationships of resistance training in healthy old adults: A systematic review and meta-analysis. *Sports Med* 45: 1693–1720, 2015.
67. Borst SE, Vincent KR, Lowenthal DT, Braith RW. Effects of resistance training on insulin-like growth factor and its binding proteins in men and women aged 60 to 85. *J Am Geriatr Soc* 50: 884–888, 2002.
68. Bosco C, Komi PV. Influence of aging on the mechanical behavior of leg extensor muscles. *Eur J Appl Physiol Occup Physiol* 45: 209–219, 1980.
69. Boshuizen HC, Stemmerik L, Westhoff MH, Hopman-Rock M. The effects of physical therapists' guidance on improvement in a strength-training program for the frail elderly. *J Aging Phys Act* 13: 5–22, 2005.
70. Boslaugh SE, Andresen EM. Peer reviewed: Correlates of physical activity for adults with disability. *Preventing Chronic Disease* 3: A78, 2006.
71. Bottaro M, Machado SN, Nogueira W, Scales R, Veloso J. Effect of high versus low-velocity resistance training on muscular fitness and functional performance in older men. *Eur J Appl Physiol* 99: 257–264, 2007.
72. Bouchard C, Rankinen T. Individual differences in response to regular physical activity. *Med Sci Sports Exerc* 33(6 Suppl): S446–S451, 2001; discussion S52–3.
73. Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: A meta-analysis of controlled clinical trials. *JAMA* 286: 1218–1227, 2001.
74. Boyle PA, Buchman AS, Wilson RS, Leurgans SE, Bennett DA. Association of muscle strength with the risk of Alzheimer disease and the rate of cognitive decline in community-dwelling older persons. *Arch Neurol* 66: 1339–1344, 2009.
75. Braith RW, Stewart KJ. Resistance exercise training: Its role in the prevention of cardiovascular disease. *Circulation* 113: 2642–2650, 2006.
76. Bray NW, Smart RR, Jakobi JM, Jones GR. Exercise prescription to reverse frailty. *Appl Physiol Nutr Metab* 41: 1112–1116, 2016.
77. Bredella MA, Gill CM, Gerweck AV, et al. Ectopic and serum lipid levels are positively associated with bone marrow fat in obesity. *Radiology* 269: 534–541, 2013.
78. Brentano MA, Cadore EL, Da Silva EM, et al. Physiological adaptations to strength and circuit training in postmenopausal women with bone loss. *J Strength Cond Res* 22: 1816–1825, 2008.
79. Brill PA, Jensen RL, Koltyn KF, et al. The feasibility of conducting a group-based progressive strength training program in residents of a multi-level care facility. *Act Adapt Aging* 22: 53–63, 1997.
80. Brill PA, Matthews M, Mason J, et al. Improving functional performance through a group-based free weight strength training program in residents of two assisted living communities. *Phys Occup Ther Geriatr* 15: 57–69, 1998.
81. Brochu M, Savage P, Lee M, et al. Effects of resistance training on physical function in older disabled women with coronary heart disease. *J Appl Physiol* (1985) 92: 672–678, 2002.
82. Brook RD, Appel LJ, Rubenfire M, et al. Beyond medications and diet: Alternative approaches to lowering blood pressure: A scientific statement from the American Heart Association. *Hypertension* 61: 1360–1383, 2013.
83. Brooks N, Layne JE, Gordon PL, et al. Strength training improves muscle quality and insulin sensitivity in Hispanic older adults with type 2 diabetes. *Int J Med Sci* 4: 19–27, 2006.
84. Brown AB, McCartney N, Sale DG. Positive adaptations to weight-lifting training in the elderly. *J Appl Physiol* (1985) 69: 1725–1733, 1990.
85. Buchman AS, Wilson RS, Boyle PA, Bienias JL, Bennett DA. Grip strength and the risk of incident Alzheimer's disease. *Neuroepidemiology* 29: 66–73, 2007.
86. Burton E, Farrier K, Lewin G, et al. Motivators and barriers for older people participating in resistance training: A systematic review. *J Aging Phys Act* 25: 311–324, 2017.
87. Bweir S, Al-Jarrah M, Almalaty AM, et al. Resistance exercise training lowers HbA1c more than aerobic training in adults with type 2 diabetes. *Diabetol Metab Syndr* 1: 27, 2009.
88. Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, et al. Multicomponent exercises including muscle power training enhance muscle mass, power output, and functional outcomes in institutionalized frail nonagenarians. *Age (Dordr)* 36: 773–785, 2014.
89. Cadore EL, Izquierdo M. New strategies for the concurrent strength-, power-, and endurance-training prescription in elderly individuals. *J Am Med Dir Assoc* 14: 623–624, 2013.
90. Cadore EL, Izquierdo M. Exercise interventions in polypathological aging patients that coexist with diabetes mellitus: Improving functional status and quality of life. *Age (Dordr)* 37: 64, 2015.
91. Cadore EL, Izquierdo M, Pinto SS, et al. Neuromuscular adaptations to concurrent training in the elderly: Effects of intrasession exercise sequence. *Age (Dordr)* 35: 891–903, 2013.
92. Cadore EL, Menger E, Teodoro JL, et al. Functional and physiological adaptations following concurrent training using sets with and without concentric failure in elderly men: A randomized clinical trial. *Exp Gerontol* 110: 182–190, 2018.
93. Cadore EL, Moneo AB, Mensat MM, et al. Positive effects of resistance training in frail elderly patients with dementia after long-term physical restraint. *Age (Dordr)* 36: 801–811, 2014.
94. Cadore EL, Pinto RS, Bottaro M, Izquierdo M. Strength and endurance training prescription in healthy and frail elderly. *Aging Dis* 5: 183–195, 2014.
95. Cadore EL, Pinto RS, Lhullier FL, et al. Physiological effects of concurrent training in elderly men. *Int J Sports Med* 31: 689–697, 2010.

96. Cadore EL, Rodríguez-Manas L, Sinclair A, Izquierdo M. Effects of different exercise interventions on risk of falls, gait ability, and balance in physically frail older adults: A systematic review. *Rejuvenation Res* 16: 105–114, 2013.
97. Calella P, Hernandez-Sanchez S, Garofalo C, Ruiz JR, Carrero JJ, Bellizzi V. Exercise training in kidney transplant recipients: A systematic review. *J Nephrol* 2019. Epub ahead of print.
98. Calle MC, Fernandez ML. Effects of resistance training on the inflammatory response. *Nutr Res Pract* 4: 259–269, 2010.
99. Campbell AJ, Borrie MJ, Spears GF. Risk factors for falls in a community-based prospective study of people 70 years and older. *J Gerontol* 44: M112–M117, 1989.
100. Campbell AJ, Buchner DM. Unstable disability and the fluctuations of frailty. *Age Ageing* 26: 315–318, 1997.
101. Campbell WW, Crim MC, Young VR, Evans WJ. Increased energy requirements and changes in body composition with resistance training in older adults. *Am J Clin Nutr* 60: 167–175, 1994.
102. Candow DG, Chilibeck PD, Facci M, Abeysekara S, Zello GA. Protein supplementation before and after resistance training in older men. *Eur J Appl Physiol* 97: 548–556, 2006.
103. Carmelli D, Reed T. Stability and change in genetic and environmental influences on hand-grip strength in older male twins. *J Appl Physiol* (1985) 89: 1879–1883, 2000.
104. Carruthers M. Testosterone deficiency syndrome: Cellular and molecular mechanism of action. *Curr Aging Sci* 6: 115–124, 2013.
105. Casas-Herrero A, Cadore EL, Zambom-Ferraresi F, et al. Functional capacity, muscle fat infiltration, power output, and cognitive impairment in institutionalized frail oldest old. *Rejuvenation Res* 16: 396–403, 2013.
106. Casas Herrero A, Cadore EL, Martínez Velilla N, Izquierdo Redin M. Physical exercise in the frail elderly: An update [in Spanish]. *Rev Esp Geriatr Gerontol* 50: 74–81, 2015.
107. Caserotti P, Aagaard P, Larsen JB, Puggaard L. Explosive heavy-resistance training in old and very old adults: Changes in rapid muscle force, strength and power. *Scand J Med Sci Sports* 18: 773–782, 2008.
108. Cassilhas RC, Antunes HK, Tufik S, de Mello MT. Mood, anxiety, and serum IGF-I in elderly men given 24 weeks of high resistance exercise. *Percept Mot Skills* 110: 265–276, 2010.
109. Castaneda C, Layne JE, Munoz-Orians L, et al. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care* 25: 2335–2341, 2002.
110. CDC—Centers for Disease Control and Prevention. Vital signs: Awareness and treatment of uncontrolled hypertension among adults — United States, 2003–2010. *MMWR Morb Mortal Wkly Rep* 61: 703–709, 2012.
111. CDC. Falls Among Older Adults: An Overview. Centers for Disease Control and Prevention. Atlanta, Georgia: National Center for Injury Prevention and Control, Division of Unintentional Injury Prevention, 2014.
112. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2017. Atlanta, GA: Centers for Disease Control and Prevention, US Department of Health and Human Services; 2017.
113. Centers for Disease C, Prevention. Self-reported falls and fall-related injuries among persons aged ≥65 years—United States, 2006. *MMWR Morb Mortal Wkly Rep* 57: 225–229, 2008.
114. Cervantes CM, Taylor WC. Physical activity interventions in adult populations with disabilities: A review. *Quest* 63: 385–410, 2011.
115. Charette SL, McEvoy L, Pyka G, et al. Muscle hypertrophy response to resistance training in older women. *J Appl Physiol* (1985) 70: 1912–1916, 1991.
116. Cheema B, Abas H, Smith B, et al. Progressive exercise for anabolism in kidney disease (PEAK): A randomized, controlled trial of resistance training during hemodialysis. *J Am Soc Nephrol* 18: 1594–1601, 2007.
117. Cheema BS, Chan D, Fahy P, Atlantis E. Effect of progressive resistance training on measures of skeletal muscle hypertrophy, muscular strength and health-related quality of life in patients with chronic kidney disease: A systematic review and meta-analysis. *Sports Med* 44: 1125–1138, 2014.
118. Chen HT, Chung YC, Chen YJ, Ho SY, Wu HJ. Effects of different types of exercise on body composition, muscle strength, and IGF-I in the elderly with sarcopenic obesity. *J Am Geriatr Soc* 65: 827–832, 2017.
119. Chen KM, Kuo CC, Chang YH, Huang HT, Cheng YY. Resistance band exercises reduce depression and behavioral problems of wheelchair-bound older adults with dementia: A cluster-randomized controlled trial. *J Am Geriatr Soc* 65: 356–363, 2017.
120. Chiu SC, Yang RS, Yang RJ, Chang SF. Effects of resistance training on body composition and functional capacity among sarcopenic obese residents in long-term care facilities: A preliminary study. *BMC Geriatr* 18: 21, 2018.
121. Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: A meta-analysis. *Arch Phys Med Rehabil* 93: 237–244, 2012.
122. Clark AL, Volterrani M, Swan JW, Coats AJ. The increased ventilatory response to exercise in chronic heart failure: Relation to pulmonary pathology. *Heart* 77: 138–146, 1997.
123. Clark BC, Manini TM. Sarcopenia 5/5 dynapenia. *J Gerontol A Biol Sci Med Sci* 63: 829–834, 2008.
124. Clark BC, Manini TM. Functional consequences of sarcopenia and dynapenia in the elderly. *Curr Opin Clin Nutr Metab Care* 13: 271–276, 2010.
125. Clemson L, Fatarone Singh MA, Bundy A, et al. Integration of balance and strength training into daily life activity to reduce rate of falls in older people (the LIFE study): Randomised parallel trial. *BMJ* 345: e4547, 2012.
126. Cohen A, Dempster DW, Recker RR, et al. Abdominal fat is associated with lower bone formation and inferior bone quality in healthy pre-menopausal women: A transilliac bone biopsy study. *J Clin Endocrinol Metab* 98: 2562–2572, 2013.
127. Cohen DD, Gomez-Arbelaez D, Camacho PA, et al. Low muscle strength is associated with metabolic risk factors in Colombian children: The ACFIES study. *PLoS One* 9: e93150, 2014.
128. Colberg SR, Sigal RJ, Yardley JE, et al. Physical activity/exercise and diabetes: A position statement of the American Diabetes Association. *Diabetes Care* 39: 2065–2079, 2016.
129. Colcombe S, Kramer AF. Fitness effects on the cognitive function of older adults: A meta-analytic study. *Psychol Sci* 14: 125–130, 2003.
130. Coleman BJ. European models of long-term care in the home and community. *Int J Health Serv* 25: 455–474, 1995.
131. Compston J, Watts N, Chapurlat R, et al. Obesity is not protective against fracture in postmenopausal women. *Am J Med* 124: 1043–1050, 2011.
132. Conlon JA, Newton RU, Tufano JJ, et al. The efficacy of periodised resistance training on neuromuscular adaptation in older adults. *Eur J Appl Physiol* 117: 1181–1194, 2017.
133. Control CfD, Prevention. Physical activity among adults with a disability—United States, 2005. *MMWR Morb Mortal Wkly Rep* 56: 1021, 2007.
134. Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 41: 1–12, 2003.
135. Coresh J, Selvin E, Stevens LA, et al. Prevalence of chronic kidney disease in the United States. *JAMA* 298: 2038–2047, 2007.
136. Cornelissen VA, Fagard RH. Effect of resistance training on resting blood pressure: A meta-analysis of randomized controlled trials. *J Hypertens* 23: 251–259, 2005.
137. Craig BW, Brown R, Everhart J. Effects of progressive resistance training on growth hormone and testosterone levels in young and elderly subjects. *Mech Ageing Dev* 49: 159–169, 1989.
138. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: Revised European consensus on definition and diagnosis. *Age Ageing* 48: 16–31, 2018.
139. Cruz-Jentoft AJ, Landi F, Schneider SM, et al. Prevalence of and interventions for sarcopenia in ageing adults: A systematic review. Report of the International Sarcopenia initiative (EWGSOP and IWGS). *Age Ageing* 43: 748–759, 2014.
140. Csapo R, Alegre LM. Effects of resistance training with moderate vs heavy loads on muscle mass and strength in the elderly: A meta-analysis. *Scand J Med Sci Sports* 26: 995–1006, 2016.
141. da Silva LXN, Teodoro JL, Menger E, et al. Repetitions to failure versus not to failure during concurrent training in healthy elderly men: A randomized clinical trial. *Exp Gerontol* 108: 18–27, 2018.
142. Dalsky GP, Stocke KS, Ehsani AA, et al. Weight-bearing exercise training and lumbar bone mineral content in postmenopausal women. *Ann Intern Med* 108: 824–828, 1988.
143. Daly M, Vidt ME, Eggebeen JD, et al. Upper extremity muscle volumes and functional strength after resistance training in older adults. *J Aging Phys Act* 21: 186–207, 2013.
144. Damush TM, Damush JG Jr. The effects of strength training on strength and health-related quality of life in older adult women. *Gerontologist* 39: 705–710, 1999.
145. Dardevet D, Sornet C, Balage M, Grizard J. Stimulation of in vitro rat muscle protein synthesis by leucine decreases with age. *J Nutr* 130: 2630–2635, 2000.
146. Davidson LE, Hudson R, Kilpatrick K, et al. Effects of exercise modality on insulin resistance and functional limitation in older adults: A randomized controlled trial. *Arch Intern Med* 169: 122–131, 2009.
147. de Boer MD, Morse CI, Thom JM, de Haan A, Narici MV. Changes in antagonist muscles' coactivation in response to strength training in older women. *J Gerontol A Biol Sci Med Sci* 62: 1022–1027, 2007.
148. De Luca CJ, Contessa P. Hierarchical control of motor units in voluntary contractions. *J Neurophysiol* 107: 178–195, 2012.
149. de Rezende LFM, Rey-Lopez JP, Matsudo VKR, do Carmo Luiz O. Sedentary behavior and health outcomes among older adults: A systematic review. *BMC Public Health* 14: 333, 2014.
150. de Vos NJ, Singh NA, Ross DA, et al. Optimal load for increasing muscle power during explosive resistance training in older adults. *J Gerontol A Biol Sci Med Sci* 60: 638–647, 2005.
151. de Vreede PL, Samson MM, van Meeteren NL, Duursma SA, Verhaar HJ. Functional-task exercise versus resistance strength exercise to improve daily function in older women: A randomized, controlled trial. *J Am Geriatr Soc* 53: 2–10, 2005.
152. de Vreede PL, van Meeteren NL, Samson MM, et al. The effect of functional tasks exercise and resistance exercise on health-related quality of life and physical activity: A randomised controlled trial. *Gerontology* 53: 12–20, 2007.
153. de Vries NM, van Ravensberg CD, Hobbelen JS, et al. Effects of physical exercise therapy on mobility, physical functioning, physical activity and quality of life in community-dwelling older adults with impaired mobility, physical disability and/or multi-morbidity: A meta-analysis. *Ageing Res Rev* 11: 136–149, 2012.
154. DeBeliso M, Harris C, Spitzer-Gibson T, Adams KJ. A comparison of periodised and fixed repetition training protocol on strength in older adults. *J Sci Med Sport* 8: 190–199, 2005.

155. Delmonico MJ, Harris TB, Visser M, et al. Longitudinal study of muscle strength, quality, and adipose tissue infiltration. *Am J Clin Nutr* 90: 1579–1585, 2009.
156. Despres JP. Body fat distribution and risk of cardiovascular disease: An update. *Circulation* 126: I301–I313, 2012.
157. Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature* 444: 881–887, 2006.
158. Despres JP, Lemieux I, Bergeron J, et al. Abdominal obesity and the metabolic syndrome: Contribution to global cardiometabolic risk. *Arterioscler Thromb Vasc Biol* 28: 1039–1049, 2008.
159. Diamanti-Kandaraki E, Dattilo M, Macut D, et al. Mechanisms in endocrinology: Aging and anti-aging: A combo-endocrinology overview. *Eur J Endocrinol* 176: R283–R308, 2017.
160. DiFrancisco-Donoghue J, Werner W, Douris PC. Comparison of once-weekly and twice-weekly strength training in older adults. *Br J Sports Med* 41: 19–22, 2007.
161. Ditroilo M, Forte R, Benelli P, Gambarara D, De Vito G. Effects of age and limb dominance on upper and lower limb muscle function in healthy males and females aged 40–80 years. *J Sports Sci* 28: 667–677, 2010.
162. Dobek JC, White KN, Gunter KB. The effect of a novel ADL-based training program on performance of activities of daily living and physical fitness. *J Aging Phys Act* 15: 13–25, 2007.
163. Doherty TJ, Brown WF. The estimated numbers and relative sizes of thenar motor units as selected by multiple point stimulation in young and older adults. *Muscle Nerve* 16: 355–366, 1993.
164. Doherty TJ, Stashuk DW, Brown WF. Determinants of mean motor unit size: Impact on estimates of motor unit number. *Muscle Nerve* 16: 1326–1331, 1993.
165. Doherty TJ, Vandervoort AA, Brown WF. Effects of ageing on the motor unit: A brief review. *Can J Appl Physiol* 18: 331–353, 1993.
166. Doherty TJ, Vandervoort AA, Taylor AW, Brown WF. Effects of motor unit losses on strength in older men and women. *J Appl Physiol* (1985) 74: 868–874, 1993.
167. Dorner T, Kranz A, Zetti-Wiedner K, Ludwig C, Rieder A, Gisinger C. The effect of structured strength and balance training on cognitive function in frail, cognitive impaired elderly long-term care residents. *Aging Clin Exp Res* 19: 400–405, 2007.
168. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 52: 377–384, 1998.
169. Duchateau J, Enoka RM. Human motor unit recordings: Origins and insight into the integrated motor system. *Brain Res* 1409: 42–61, 2011.
170. Duchowny KA, Peterson MD, Clarke PJ. Cut points for clinical muscle weakness among older Americans. *Am J Prev Med* 53: 63–69, 2017.
171. Dunstan DW, Daly RM, Owen N, et al. High-intensity resistance training improves glycaemic control in older patients with type 2 diabetes. *Diabetes Care* 25: 1729–1736, 2002.
172. Evans JW. Periodized resistance training for enhancing skeletal muscle hypertrophy and strength: A mini-review. *Front Physiol* 10: 13, 2019.
173. Evans W. Functional and metabolic consequences of sarcopenia. *J Nutr* 127(5 Suppl): 998S–1003S, 1997.
174. Evans WJ. Effects of exercise on senescent muscle. *Clin Orthop Relat Res* (403 Suppl): S211–S220, 2002.
175. Fabbri E, An Y, Gonzalez-Freire M, et al. Bioavailable testosterone declines over a wide age spectrum in men and women from the Baltimore longitudinal study of aging. *J Gerontol A Biol Sci Med Sci* 71: 1202–1209, 2016.
176. Fagard RH. Epidemiology of hypertension in the elderly. *Am J Geriatr Cardiol* 11: 23–28, 2002.
177. Fatouros IG, Kambas A, Katrabasas I, et al. Resistance training and detraining effects on flexibility performance in the elderly are intensity-dependent. *J Strength Cond Res* 20: 634–642, 2006.
178. Fatouros IG, Taxildaris K, Tokmakidis SP, et al. The effects of strength training, cardiovascular training and their combination on flexibility of inactive older adults. *Int J Sports Med* 23: 112–119, 2002.
179. Featherstone JF, Holly RG, Amsterdam EA. Physiologic responses to weight lifting in coronary artery disease. *Am J Cardiol* 71: 287–292, 1993.
180. Ferrari R, Krueger LF, Cadore EL, et al. Efficiency of twice weekly concurrent training in trained elderly men. *Exp Gerontol* 48: 1236–1242, 2013.
181. Ferri A, Scaglioni G, Pousson M, Capodaglio P, Van Hoecke J, Narici MV. Strength and power changes of the human plantar flexors and knee extensors in response to resistance training in old age. *Acta Physiol Scand* 177: 69–78, 2003.
182. Fiatarone MA, Kehayias JJ, Roberts SB, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *New Engl J Med* 330: 1769–1775, 1994.
183. Fiatarone MA, Marks EC, Ryan ND, Meredith CN, Lipsitz LA, Evans WJ. High-intensity strength training in nonagenarians. Effects on skeletal muscle. *JAMA* 263: 3029–3034, 1990.
184. Fiatarone MA, O'Neill EF, Ryan ND, et al. Exercise training and nutritional supplementation for physical frailty in very elderly people. *N Engl J Med* 330: 1769–1775, 1994.
185. Fiatarone Singh MA, Gates N, Saigal N, et al. The study of mental and resistance training (SMART) study—resistance training and/or cognitive training in mild cognitive impairment: A randomized, double-blind, double-sham controlled trial. *J Am Med Dir Assoc* 15: 873–880, 2014.
186. Fielding RA, LeBrasseur NK, Cuoco A, Bean J, Mizer K, Fiatarone Singh MA. High-velocity resistance training increases skeletal muscle peak power in older women. *J Am Geriatr Soc* 50: 655–662, 2002.
187. Fisher NM, Pendergast DR, Calkins E. Muscle rehabilitation in impaired elderly nursing home residents. *Arch Phys Med Rehabil* 72: 181–185, 1991.
188. Fleg JL, Cooper LS, Borlaug BA, et al. Exercise training as therapy for heart failure: Current status and future directions. *Circ Heart Fail* 8: 209–220, 2015.
189. Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. Trends in obesity among adults in the United States, 2005 to 2014. *JAMA* 315: 2284–2291, 2016.
190. Foldvari M, Clark M, Laviolette LC, et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci* 55: M192–M199, 2000.
191. Forman-Hoffman VL, Ault KL, Anderson WL, et al. Disability status, mortality, and leading causes of death in the United States community population. *Med Care* 53: 346, 2015.
192. Fragala MS, Alley DE, Shardell MD, et al. Comparison of handgrip and leg extension strength in predicting slow gait speed in older adults. *J Am Geriatr Soc* 64: 144–150, 2016.
193. Franceschi C, Campisi J. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol A Biol Sci Med Sci* 69(Suppl 1): S4–S9, 2014.
194. Frederiksen H, Gaist D, Petersen HC, et al. Hand grip strength: A phenotype suitable for identifying genetic variants affecting mid- and late-life physical functioning. *Genet Epidemiol* 23: 110–122, 2002.
195. Freiburger E, Haberle L, Spirduso WW, Zijlstra GA. Long-term effects of three multi-component exercise interventions on physical performance and fall-related psychological outcomes in community-dwelling older adults: A randomized controlled trial. *J Am Geriatr Soc* 60: 437–446, 2012.
196. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: Evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 56: M146–M156, 2001.
197. Frontera WR. What is it about old muscles? *J Physiol* 595: 4581–4582, 2017.
198. Frontera WR, Bean JF, Damiano D, et al. Rehabilitation research at the National Institutes of Health: Moving the field forward (executive summary). *Arch Phys Med Rehabil* 98: 795–803, 2017.
199. Frontera WR, Hughes VA, Fielding RA, et al. Aging of skeletal muscle: A 12-yr longitudinal study. *J Appl Physiol* (1985) 88: 1321–1326, 2000.
200. Frontera WR, Hughes VA, Lutz KJ, Evans WJ. A cross-sectional study of muscle strength and mass in 45- to 78-yr-old men and women. *J Appl Physiol* (1985) 71: 644–650, 1991.
201. Frontera WR, Meredith CN, O'Reilly KP, Knuttigen HG, Evans WJ. Strength conditioning in older men: Skeletal muscle hypertrophy and improved function. *J Appl Physiol* (1985) 64: 1038–1044, 1988.
202. Frontera WR, Suh D, Krivickas LS, Hughes VA, Goldstein R, Roubenoff R. Skeletal muscle fiber quality in older men and women. *Am J Physiol Cell Physiol* 279: C611–C618, 2000.
203. Fyfe JJ, Bishop DJ, Stepto NK. Interference between concurrent resistance and endurance exercise: Molecular bases and the role of individual training variables. *Sports Med* 44: 743–762, 2014.
204. Gadelha AB, Paiva FM, Gauche R, de Oliveira RJ, Lima RM. Effects of resistance training on sarcopenic obesity index in older women: A randomized controlled trial. *Arch Gerontol Geriatr* 65: 168–173, 2016.
205. Gaffey AE, Bergeman CS, Clark LA, Wirth MM. Aging and the HPA axis: Stress and resilience in older adults. *Neurosci Biobehav Rev* 68: 928–945, 2016.
206. Gallagher D, Kelley DE, Yim JE, et al. Adipose tissue distribution is different in type 2 diabetes. *Am J Clin Nutr* 89: 807–814, 2009.
207. Gallagher D, Visser M, De Meersman RE, et al. Appendicular skeletal muscle mass: Effects of age, gender, and ethnicity. *J Appl Physiol* (1985) 83: 229–239, 1997.
208. Galvao DA, Newton RU, Taaffe DR. Anabolic responses to resistance training in older men and women: A brief review. *J Aging Phys Act* 13: 343–358, 2005.
209. Galvao DA, Taaffe DR. Resistance exercise dosage in older adults: Single- versus multiset effects on physical performance and body composition. *J Am Geriatr Soc* 53: 2090–2097, 2005.
210. Garatachea N, Lucia A. Genes and the ageing muscle: A review on genetic association studies. *Age (Dordr)* 35: 207–233, 2013.
211. Garcia-Garcia FJ, Gutierrez Avila G, Alfaro-Acha A, et al. The prevalence of frailty syndrome in an older population from Spain. The Toledo Study for Healthy Aging. *J Nutr Health Aging* 15: 852–856, 2011.
212. Gassen NC, Chrousos GP, Binder EB, Zannas AS. Life stress, glucocorticoid signaling, and the aging epigenome: Implications for aging-related diseases. *Neurosci Biobehav Rev* 74: 356–365, 2017.
213. Gatti R, De Palo EF, Antonelli G, Spinella P. IGF-1/IGFBP system: Metabolism outline and physical exercise. *J Endocrinol Invest* 35: 699–707, 2012.
214. Gauche R, Lima RM, Myers J, et al. Blood pressure reactivity to mental stress is attenuated following resistance exercise in older hypertensive women. *Clin Interv Aging* 12: 793–803, 2017.
215. Gell NM, Wallace RB, LaCroix AZ, Mroz TM, Patel KV. Mobility device use in older adults and incidence of falls and worry about falling: Findings from the 2011–2012 national health and aging trends study. *J Am Geriatr Soc* 63: 853–859, 2015.

216. Gerstner GR, Thompson BJ, Rosenberg JG, et al. Neural and muscular contributions to the age-related reductions in rapid strength. *Med Sci Sports Exerc* 49: 1331–1339, 2017.
217. Ghadieh AS, Saab B. Evidence for exercise training in the management of hypertension in adults. *Can Fam Physician* 61: 233–239, 2015.
218. Giangregorio LM, Papaioannou A, Macintyre NJ, et al. Too fit to fracture: Exercise recommendations for individuals with osteoporosis or osteoporotic vertebral fracture. *Osteoporos Int* 25: 821–835, 2014.
219. Gilbert KL, Stokes KA, Hall GM, Thompson D. Growth hormone responses to 3 different exercise bouts in 18- to 25- and 40- to 50-year-old men. *Appl Physiol Nutr Metab* 33: 706–712, 2008.
220. Gine ' -Garriga M, Guerra M, Page ' s E, et al. The effect of functional circuit training on physical frailty in frail older adults: A randomized controlled trial. *J Aging Phys Act* 18: 401–424, 2010.
221. Gonzalez AM, Mangine GT, Fragala MS, et al. Resistance training improves single leg stance performance in older adults. *Aging Clin Exp Res* 26: 89–92, 2014.
222. Goodpaster BH, Carlson CL, Visser M, et al. Attenuation of skeletal muscle and strength in the elderly: The Health ABC Study. *J Appl Physiol* 90: 2157–2165, 2001.
223. Goodpaster BH, Chomentowski P, Ward BK, et al. Effects of physical activity on strength and skeletal muscle fat infiltration in older adults: A randomized controlled trial. *J Appl Physiol* (1985) 105: 1498–1503, 2008.
224. Goodpaster BH, Kelley DE, Thaete FL, He J, Ross R. Skeletal muscle attenuation determined by computed tomography is associated with skeletal muscle lipid content. *J Appl Physiol* 89: 104–110, 2000.
225. Goodpaster BH, Park SW, Harris TB, et al. The loss of skeletal muscle strength, mass, and quality in older adults: The health, aging and body composition study. *J Gerontol A Biol Sci Med Sci* 61: 1059–1064, 2006.
226. Goodpaster BH, Thaete FL, Kelley DE. Thigh adipose tissue distribution is associated with insulin resistance in obesity and in type 2 diabetes mellitus. *Am J Clin Nutr* 71: 885–892, 2000.
227. Gordon NF, Kohl HW III, Pollock ML, et al. Cardiovascular safety of maximal strength testing in healthy adults. *Am J Cardiol* 76: 851–853, 1995.
228. Gordon SE, Kraemer WJ, Looney DP, et al. The influence of age and exercise modality on growth hormone bioactivity in women. *Growth Horm IGF Res* 24: 95–103, 2014.
229. Gordon SE, Kraemer WJ, Vos NH, Lynch JM, Knuttgen HG. Effect of acid-base balance on the growth hormone response to acute high-intensity cycle exercise. *J Appl Physiol* (1985) 76: 821–829, 1994.
230. Gorostiaga EM, Navarro-Amezqueta I, Calbet JA, et al. Energy metabolism during repeated sets of leg press exercise leading to failure or not. *PLoS One* 7: e40621, 2012.
231. Gower BA, Casazza K. Divergent effects of obesity on bone health. *J Clin Densitom* 16: 450–454, 2013.
232. Granacher U, Gruber M, Gollhofer A. Resistance training and neuromuscular performance in seniors. *Int J Sports Med* 30: 652–657, 2009.
233. Gregg EW, Sattar N, Ali MK. The changing face of diabetes complications. *Lancet Diabetes Endocrinol* 4: 537–547, 2016.
234. Gregg EW, Sorlie P, Paulose-Ram R, et al. Prevalence of lower-extremity disease in the US adult population .540 years of age with and without diabetes: 1999–2000 national health and nutrition examination survey. *Diabetes Care* 27: 1591–1597, 2004.
235. Gregg EW, Zhuo X, Cheng YJ, Albright AL, Narayan KMV, Thompson TJ. Trends in lifetime risk and years of life lost due to diabetes in the USA, 1985–2011: A modelling study. *Lancet Diabetes Endocrinol* 2: 867–874, 2014.
236. Greiwe JS, Cheng B, Rubin DC, Yarasheski KE, Semenkovich CF. Resistance exercise decreases skeletal muscle tumor necrosis factor α in frail elderly humans. *FASEB J* 15: 475–482, 2001.
237. Grontved A, Rimm EB, Willett WC, Andersen LB, Hu FB. A prospective study of weight training and risk of type 2 diabetes mellitus in men. *Arch Intern Med* 172: 1306–1312, 2012.
238. Hakkinen K, Hakkinen A. Neuromuscular adaptations during intensive strength training in middle-aged and elderly males and females. *Electromyogr Clin Neurophysiol* 35: 137–147, 1995.
239. Hakkinen K, Kallinen M, Izquierdo M, et al. Changes in agonist-antagonist EMG, muscle CSA, and force during strength training in middle-aged and older people. *J Appl Physiol* (1985) 84: 1341–1349, 1998.
240. Hakkinen K, Kallinen M, Linnamo V, et al. Neuromuscular adaptations during bilateral versus unilateral strength training in middle-aged and elderly men and women. *Acta Physiol Scand* 158: 77–88, 1996.
241. Hakkinen K, Kraemer WJ, Newton RU, Alen M. Changes in electromyographic activity, muscle fiber and force production characteristics during heavy resistance/power strength training in middle-aged and older men and women. *Acta Physiol Scand* 171: 51–62, 2001.
242. Hakkinen K, Kraemer WJ, Pakarinen A, et al. Effects of heavy resistance/power training on maximal strength, muscle morphology, and hormonal response patterns in 60–75-year-old men and women. *Can J Appl Physiol* 27: 213–231, 2002.
243. Hakkinen K, Newton RU, Gordon SE, et al. Changes in muscle morphology, electromyographic activity, and force production characteristics during progressive strength training in young and older men. *J Gerontol A Biol Sci Med Sci* 53: B415–B423, 1998.
244. Hakkinen K, Pakarinen A. Serum hormones and strength development during strength training in middle-aged and elderly males and females. *Acta Physiol Scand* 150: 211–219, 1994.
245. Hakkinen K, Pakarinen A, Kraemer WJ, et al. Selective muscle hypertrophy, changes in EMG and force, and serum hormones during strength training in older women. *J Appl Physiol* (1985) 91: 569–580, 2001.
246. Hakkinen K, Pakarinen A, Kraemer WJ, Newton RU, Alen M. Basal concentrations and acute responses of serum hormones and strength development during heavy resistance training in middle-aged and elderly men and women. *J Gerontol A Biol Sci Med Sci* 55: B95–B105, 2000.
247. Hakkinen K, Pakarinen A, Newton RU, Kraemer WJ. Acute hormone responses to heavy resistance lower and upper extremity exercise in young versus old men. *Eur J Appl Physiol Occup Physiol* 77: 312–319, 1998.
248. Hakkinen K, Pastinen UM, Karsikas R, Linnamo V. Neuromuscular performance in voluntary bilateral and unilateral contraction and during electrical stimulation in men at different ages. *Eur J Appl Physiol Occup Physiol* 70: 518–527, 1995.
249. Hameed M, Lange KH, Andersen JL, et al. The effect of recombinant human growth hormone and resistance training on IGF-1 mRNA expression in the muscles of elderly men. *J Physiol* 555(Pt 1): 231–240, 2004.
250. Hameed M, Orrell RW, Cobbold M, Goldspink G, Harridge SD. Expression of IGF-1 splice variants in young and old human skeletal muscle after high resistance exercise. *J Physiol* 547(Pt 1): 247–254, 2003.
251. Harridge SD, Kryger A, Stensgaard A. Knee extensor strength, activation, and size in very elderly people following strength training. *Muscle Nerve* 22: 831–839, 1999.
252. Hassan BH, Hewitt J, Keogh JW, et al. Impact of resistance training on sarcopenia in nursing care facilities: A pilot study. *Geriatr Nurs* 37: 116–121, 2016.
253. Hasten DL, Pak-Loduca J, Obert KA, Yarasheski KE. Resistance exercise acutely increases MHC and mixed muscle protein synthesis rates in 78–84 and 23–32 yr olds. *Am J Physiol Endocrinol Metab* 278: E620–E626, 2000.
254. Haub MD, Wells AM, Tarnopolsky MA, Campbell WW. Effect of protein source on resistance-training-induced changes in body composition and muscle size in older men. *Am J Clin Nutr* 76: 511–517, 2002.
255. Hauer K, Rost B, Rutschle K, et al. Exercise training for rehabilitation and secondary prevention of falls in geriatric patients with a history of injurious falls. *J Am Geriatr Soc* 49: 10–20, 2001.
256. Hauer K, Schwenk M, Zieschang T, Essig M, Becker C, Oster P. Physical training improves motor performance in people with dementia: A randomized controlled trial. *J Am Geriatr Soc* 60: 8–15, 2012.
257. Haykowsky M, Humen D, Teo K, et al. Effects of 16 weeks of resistance training on left ventricular morphology and systolic function in healthy men .60 years of age. *Am J Cardiol* 85: 1002–1006, 2000.
258. Haykowsky M, McGavock J, Vonder Muhll I, et al. Effect of exercise training on peak aerobic power, left ventricular morphology, and muscle strength in healthy older women. *J Gerontol A Biol Sci Med Sci* 60: 307–311, 2005.
259. Hazell T, Kenno K, Jakobi J. Functional benefit of power training for older adults. *J Aging Phys Act* 15: 349–359, 2007.
260. Heffernan KS, Yoon ES, Sharman JE, et al. Resistance exercise training reduces arterial reservoir pressure in older adults with prehypertension and hypertension. *Hypertens Res* 36: 422–427, 2013.
261. Hennessey JV, Chromiak JA, DellaVentura S, et al. Growth hormone administration and exercise effects on muscle fiber type and diameter in moderately frail older people. *J Am Geriatr Soc* 49: 852–858, 2001.
262. Henwood TR, Riek S, Taaffe DR. Strength versus muscle power-specific resistance training in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 63: 83–91, 2008.
263. Henwood TR, Taaffe DR. Improved physical performance in older adults undertaking a short-term programme of high-velocity resistance training. *Gerontology* 51: 108–115, 2005.
264. Henwood TR, Taaffe DR. Short-term resistance training and the older adult: The effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging* 26: 305–313, 2006.
265. Heyn P, Abreu BC, Ottenbacher KJ. The effects of exercise training on elderly persons with cognitive impairment and dementia: A meta-analysis. *Arch Phys Med Rehabil* 85: 1694–1704, 2004.
266. Heyn PC, Johnson KE, Kramer AF. Endurance and strength training outcomes on cognitively impaired and cognitively intact older adults: A meta-analysis. *J Nutr Health Aging* 12: 401–409, 2008.
267. Hickson RC. Interference of strength development by simultaneously training for strength and endurance. *Eur J Appl Physiol Occup Physiol* 45: 255–263, 1980.
268. Hoffmann TC, Maher CG, Briffa T, et al. Prescribing exercise interventions for patients with chronic conditions. *CMAJ* 188: 510–518, 2016.
269. Hofmann M, Schober-Halper B, Oesen S, et al. Effects of elastic band resistance training and nutritional supplementation on muscle quality and circulating muscle growth and degradation factors of institutionalized elderly women: The Vienna Active Ageing Study (VAAS). *Eur J Appl Physiol* 116: 885–897, 2016.
270. Hollings M, Mavros Y, Freeston J, Fatarone Singh M. The effect of progressive resistance training on aerobic fitness and strength in adults with coronary heart disease: A systematic review and meta-analysis of randomised controlled trials. *Eur J Prev Cardiol* 24: 1242–1259, 2017.

271. Holviala JH, Sallinen JM, Kraemer WJ, Alen MJ, Hakkinen KK. Effects of strength training on muscle strength characteristics, functional capabilities, and balance in middle-aged and older women. *J Strength Cond Res* 20: 336–344, 2006.
272. Hortobagyi T, Tunnel D, Moody J, Beam S, DeVita P. Low- or high- intensity strength training partially restores impaired quadriceps force accuracy and steadiness in aged adults. *J Gerontol A Biol Sci Med Sci* 56: B38–B47, 2001.
273. Hovanec N, Sawant A, Overend TJ, Petrella RJ, Vandervoort AA. Resistance training and older adults with type 2 diabetes mellitus: Strength of the evidence. *J Aging Res* 2012: 284635, 2012.
274. Hruska KV, Hicks AL, McCartney N. Training for muscle power in older adults: Effects on functional abilities. *Can J Appl Physiol* 28: 178–189, 2003.
275. Hubal MJ, Gordish-Dressman H, Thompson PD, et al. Variability in muscle size and strength gain after unilateral resistance training. *Med Sci Sports Exerc* 37: 964–972, 2005.
276. Hulmi JJ, Ahtiainen JP, Selanne H, et al. Androgen receptors and testosterone in men—Effects of protein ingestion, resistance exercise and fiber type. *J Steroid Biochem Mol Biol* 110: 130–137, 2008.
277. Hunter GR, McCarthy JP, Bamman MM. Effects of resistance training on older adults. *Sports Med* 34: 329–348, 2004.
278. Hunter GR, Wetzstein CJ, Fields DA, Brown A, Bamman MM. Resistance training increases total energy expenditure and free-living physical activity in older adults. *J Appl Physiol* (1985) 89: 977–984, 2000.
279. Hunter GR, Wetzstein CJ, McLafferty CL Jr, et al. High-resistance versus variable-resistance training in older adults. *Med Sci Sports Exerc* 33: 1759–1764, 2001.
280. Hunter SK, Pereira HM, Keenan KG. The aging neuromuscular system and motor performance. *J Appl Physiol* (1985) 121: 982–995, 2016.
281. Hurlbut DE, Lott ME, Ryan AS, et al. Does age, sex, or ACE genotype affect glucose and insulin responses to strength training? *J Appl Physiol* (1985) 92: 643–650, 2002.
282. Hurley BF, Hanson ED, Sheaff AK. Strength training as a countermeasure to aging muscle and chronic disease. *Sports Med* 41: 289–306, 2011.
283. Hyatt RH, Whitelaw MN, Bhat A, Scott S, Maxwell JD. Association of muscle strength with functional status of elderly people. *Age Ageing* 19: 330–336, 1990.
284. Ibanez J, Izquierdo M, Arguëlles I, et al. Twice-weekly progressive resistance training decreases abdominal fat and improves insulin sensitivity in older men with type 2 diabetes. *Diabetes Care* 28: 662–667, 2005.
285. Ice GH. Daily life in a nursing home Has it changed in 25 years? *J Aging Stud* 16: 345, 2002.
286. Igwebuike A, Irving BA, Bigelow ML, et al. Lack of dehydroepiandrosterone effect on a combined endurance and resistance exercise program in postmenopausal women. *J Clin Endocrinol Metab* 93: 534–538, 2008.
287. Ivey FM, Roth SM, Ferrell RE, et al. Effects of age, gender, and myostatin genotype on the hypertrophic response to heavy resistance strength training. *J Gerontol A Biol Sci Med Sci* 55: M641–M648, 2000.
288. Ivey FM, Tracy BL, Lemmer JT, et al. Effects of strength training and detraining on muscle quality: Age and gender comparisons. *J Gerontol A Biol Sci Med Sci* 55: B152–B157, 2000; discussion B8–9.
289. Izquierdo M, Hakkinen K, Anton A, et al. Maximal strength and power, endurance performance, and serum hormones in middle-aged and elderly men. *Med Sci Sports Exerc* 33: 1577–1587, 2001.
290. Izquierdo M, Hakkinen K, Ibanez J, et al. Effects of strength training on submaximal and maximal endurance performance capacity in middle-aged and older men. *J Strength Cond Res* 17: 129–139, 2003.
291. Izquierdo M, Hakkinen K, Ibanez J, et al. Effects of strength training on muscle power and serum hormones in middle-aged and older men. *J Appl Physiol* (1985) 90: 1497–1507, 2001.
292. Izquierdo M, Ibanez J, Gorostiaga E, et al. Maximal strength and power characteristics in isometric and dynamic actions of the upper and lower extremities in middle-aged and older men. *Acta Physiol Scand* 167: 57–68, 1999.
293. Izquierdo M, Ibanez J, Kraemer WJ, Larrion JL, Gorostiaga EM. Once weekly combined resistance and cardiovascular training in healthy older men. *Med Sci Sports Exerc* 36: 435–443, 2004.
294. Izumi A, Kitamura M, Izawa KP. Effects of exercise training on delaying disease progression in patients with chronic kidney disease: A review of the literature. *Rev Recent Clin Trials* 11: 333–341, 2016.
295. Jafarzadeh SR, Felson DT. Updated estimates suggest a much higher prevalence of arthritis in US adults than previous ones. *Arthritis Rheumatol* 70: 185–192, 2017.
296. Jakobi JM, Rice CL. Voluntary muscle activation varies with age and muscle group. *J Appl Physiol* (1985) 93: 457–462, 2002.
297. Jan MH, Lin JJ, Liao JJ, Lin YF, Lin DH. Investigation of clinical effects of high- and low-resistance training for patients with knee osteoarthritis: A randomized controlled trial. *Phys Ther* 88: 427–436, 2008.
298. Janssen I, Heymsfield SB, Wang ZM, Ross R. Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J Appl Physiol* (1985) 89: 81–88, 2000.
299. Janssen JA. Impact of physical exercise on endocrine aging. *Front Horm Res* 47: 68–81, 2016.
300. Johansen KL, Painter P. Exercise in individuals with CKD. *Am J Kidney Dis* 59: 126–134, 2012.
301. Johnen B, Schott N. Feasibility of a machine vs free weight strength training program and its effects on physical performance in nursing home residents: A pilot study. *Aging Clin Exp Res* 30: 819–828, 2018.
302. Johnson RJ, Rodríguez-Iturbe B, Roncal-Jimenez C, et al. Hyperosmolarity drives hypertension and CKD—water and salt revisited. *Nat Rev Nephrol* 10: 415–420, 2014.
303. Johnson Stoklossa CA, Sharma AM, Forhan M, et al. Prevalence of sarcopenic obesity in adults with class III/III obesity using different diagnostic criteria. *J Nutr Metab* 2017: 7307618, 2017.
304. Johnston AP, De Lisio M, Parise G. Resistance training, sarcopenia, and the mitochondrial theory of aging. *Appl Physiol Nutr Metab* 33: 191–199, 2008.
305. Jordan JM, Helmick CG, Renner JB, et al. Prevalence of knee symptoms and radiographic and symptomatic knee osteoarthritis in African Americans and Caucasians: The Johnston county osteoarthritis project. *J Rheumatol* 34: 172–180, 2007.
306. Jordan JM, Helmick CG, Renner JB, et al. Prevalence of hip symptoms and radiographic and symptomatic hip osteoarthritis in African Americans and Caucasians: The Johnston county osteoarthritis project. *J Rheumatol* 36: 809–815, 2009.
307. Joseph LJ, Davey SL, Evans WJ, Campbell WW. Differential effect of resistance training on the body composition and lipoprotein-lipid profile in older men and women. *Metabolism* 48: 1474–1480, 1999.
308. Judge JO, Whipple RH, Wolfson LI. Effects of resistive and balance exercises on isokinetic strength in older persons. *J Am Geriatr Soc* 42: 937–946, 1994.
309. Kalapotharakos V, Smilios I, Parlavatzas A, Tokmakidis SP. The effect of moderate resistance strength training and detraining on muscle strength and power in older men. *J Geriatr Phys Ther* 30: 109–113, 2007.
310. Kalapotharakos VI, Michalopoulou M, Godolias G, et al. The effects of high- and moderate-resistance training on muscle function in the elderly. *J Aging Phys Act* 12: 131–143, 2004.
311. Kalinkovich A, Livshits G. Sarcopenic obesity or obese sarcopenia: A cross talk between age-associated adipose tissue and skeletal muscle inflammation as a main mechanism of the pathogenesis. *Ageing Res Rev* 35: 200–221, 2017.
312. Kalyani RR, Metter EJ, Egan J, Golden SH, Ferrucci L. Hyperglycemia predicts persistently lower muscle strength with aging. *Diabetes Care* 38: 82–90, 2015.
313. Kane RL, Butler M, Fink HA, et al. Interventions to Prevent Age-Related Cognitive Decline, Mild Cognitive Impairment, and Clinical Alzheimer's-Type Dementia. Rockville, MD: Agency for Healthcare Research and Quality (US), 2017.
314. Karamanidis K, Arampatzis A. Mechanical and morphological properties of human quadriceps femoris and triceps surae muscle-tendon unit in relation to aging and running. *J Biomech* 39: 406–417, 2006.
315. Karinkanta S, Heinonen A, Sievanen H, et al. A multi-component exercise regimen to prevent functional decline and bone fragility in home-dwelling elderly women: Randomized, controlled trial. *Osteoporos Int* 18: 453–462, 2007.
316. Karinkanta S, Kannus P, Uusi-Rasi K, Heinonen A, Sievanen H. Combined resistance and balance-jumping exercise reduces older women's injurious falls and fractures: 5-year follow-up study. *Age Ageing* 44: 784–789, 2015.
317. Kekalainen T, Kokko K, Sipilä S, Walker S. Effects of a 9-month resistance training intervention on quality of life, sense of coherence, and depressive symptoms in older adults: Randomized controlled trial. *Qual Life Res* 27: 455–465, 2017.
318. Kelley GA, Kelley KS. Progressive resistance exercise and resting blood pressure: A meta-analysis of randomized controlled trials. *Hypertension* 35: 838–843, 2000.
319. Kelsey TW, Li LQ, Mitchell RT, et al. A validated age-related normative model for male total testosterone shows increasing variance but no decline after age 40 years. *PLoS One* 9: e109346, 2014.
320. Kendrick D, Kumar A, Carpenter H, et al. Exercise for reducing fear of falling in older people living in the community. *Cochrane Database Syst Rev* 11: CD009848, 2014.
321. Kennis E, Verschueren SM, Bogaerts A, et al. Long-term impact of strength training on muscle strength characteristics in older adults. *Arch Phys Med Rehabil* 94: 2054–2060, 2013.
322. Kessenich CR. Health-related quality of life in osteoporosis. *J Clin Densitom* 1: 27–31, 1998.
323. Keysor JJ. Does late-life physical activity or exercise prevent or minimize disability? A critical review of the scientific evidence. *Am J Prev Med* 25(3 Suppl 2): 129–136, 2003.
324. Keysor JJ, Jette AM. Have we oversold the benefit of late-life exercise? *J Gerontol A Biol Sci Med Sci* 56: M412–M423, 2001.
325. Kimball SR, Jefferson LS. Control of protein synthesis by amino acid availability. *Curr Opin Clin Nutr Metab Care* 5: 63–67, 2002.
326. King LK, Birmingham TB, Kean CO, Jones IC, Bryant DM, Giffin JR. Resistance training for medial compartment knee osteoarthritis and malalignment. *Med Sci Sports Exerc* 40: 1376–1384, 2008.
327. Kinne S, Patrick DL, Doyle DL. Prevalence of secondary conditions among people with disabilities. *Am J Public Health* 94: 443–445, 2004.
328. Kitwood T. The dialectics of dementia: With particular reference to Alzheimer's disease. *Ageing Soc* 10: 177–196, 1990.
329. Klass M, Baudry S, Duchateau J. Voluntary activation during maximal contraction with advancing age: A brief review. *Eur J Appl Physiol* 100: 543–551, 2007.
330. Klass M, Baudry S, Duchateau J. Age-related decline in rate of torque development is accompanied by lower maximal motor unit discharge frequency during fast contractions. *J Appl Physiol* (1985) 104: 739–746, 2008.
331. Klein CS, Ivanova TD, Rice CL, Garland SJ. Motor unit discharge rate following twitch

- potentiation in human triceps brachii muscle. *Neurosci Lett* 316: 153–156, 2001.
332. Klitgaard H, Mantoni M, Schiaffino S, et al. Function, morphology and protein expression of ageing skeletal muscle: A cross-sectional study of elderly men with different training backgrounds. *Acta Physiol Scand* 140: 41–54, 1990.
 333. Knight CA, Kamen G. Adaptations in muscular activation of the knee extensor muscles with strength training in young and older adults. *J Electromyogr Kinesiol* 11: 405–412, 2001.
 334. Kolber MJ, Beekhuizen KS, Cheng MS, Hellman MA. Shoulder injuries attributed to resistance training: A brief review. *J Strength Cond Res* 24: 1696–1704, 2010.
 335. Korpelainen R, Keinanen-Kiukkaanniemi S, Heikkinen J, Vaananen K, Korpelainen J. Effect of impact exercise on bone mineral density in elderly women with low BMD: A population-based randomized controlled 30-month intervention. *Osteoporos Int* 17: 109–118, 2006.
 336. Kortebein P, Symons TB, Ferrando A, et al. Functional impact of 10 days of bed rest in healthy older adults. *J Gerontol A Biol Sci Med Sci* 63: 1076–1081, 2008.
 337. Kosek DJ, Kim JS, Petrella JK, Cross JM, Bamman MM. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J Appl Physiol* (1985) 101: 531–544, 2006.
 338. Kraemer WJ. Endocrine responses to resistance exercise. *Med Sci Sports Exerc* 20(5 Suppl): S152–S157, 1988.
 339. Kraemer WJ, Deschenes MR, Fleck SJ. Physiological adaptations to resistance exercise. Implications for athletic conditioning. *Sports Med* 6: 246–256, 1988.
 340. Kraemer WJ, Dunn-Lewis C, Comstock BA, Thomas GA, Clark JE, Nindl BC. Growth hormone, exercise, and athletic performance: A continued evolution of complexity. *Curr Sports Med Rep* 9: 242–252, 2010.
 341. Kraemer WJ, Fragala MS. Personalize it: Program design in resistance training. *ACSM Health Fitness J* 10: 7–17, 2006.
 342. Kraemer WJ, Hakkinen K, Newton RU, et al. Effects of heavy-resistance training on hormonal response patterns in younger vs. older men. *J Appl Physiol* (1985) 87: 982–992, 1999.
 343. Kraemer WJ, Kennett MJ, Mastro AM, et al. Bioactive growth hormone in older men and women: It's relationship to immune markers and healthspan. *Growth Horm IGF Res* 34: 45–54, 2017.
 344. Kraemer WJ, Nindl BC, Marx JO, et al. Chronic resistance training in women potentiates growth hormone in vivo bioactivity: Characterization of molecular mass variants. *Am J Physiol Endocrinol Metab* 291: E1177–E1187, 2006.
 345. Kraemer WJ, Patton JF, Gordon SE, et al. Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations. *J Appl Physiol* (1985) 78: 976–989, 1995.
 346. Kraemer WJ, Ratamess NA. Hormonal responses and adaptations to resistance exercise and training. *Sports Med* 35: 339–361, 2005.
 347. Kraemer WJ, Ratamess NA, Nindl BC. Recovery responses of testosterone, growth hormone, and IGF-I after resistance exercise. *J Appl Physiol* (1985) 122: 549–558, 2017.
 348. Kraemer WJ, Spiering BA, Volek JS, et al. Androgenic responses to resistance exercise: Effects of feeding and L-carnitine. *Med Sci Sports Exerc* 38: 1288–1296, 2006.
 349. Krebs DE, Scarborough DM, McGibbon CA. Functional vs. strength training in disabled elderly outpatients. *Am J Phys Med Rehabil* 86: 93–103, 2007.
 350. Krist L, Dimeo F, Keil T. Can progressive resistance training twice a week improve mobility, muscle strength, and quality of life in very elderly nursing-home residents with impaired mobility? A pilot study. *Clin Interv Aging* 8: 443–448, 2013.
 351. Krops LA, Hols DH, Folkertsma N, et al. Requirements on a community-based intervention for stimulating physical activity in physically disabled people: A focus group study amongst experts. *Disabil Rehabil* 40: 2400–2407, 2018.
 352. Kryger AI, Andersen JL. Resistance training in the oldest old: Consequences for muscle strength, fiber types, fiber size, and MHC isoforms. *Scand J Med Sci Sports* 17: 422–430, 2007.
 353. Ku PW, Fox KR, Gardiner PA, Chen LJ. Late-life exercise and difficulty with activities of daily living: An 8-year nationwide follow-up study in Taiwan. *Ann Behav Med* 50: 237–246, 2016.
 354. Lacroix A, Hortobagyi T, Beurskens R, Granacher U. Effects of supervised vs. unsupervised training programs on balance and muscle strength in older adults: A systematic review and meta-analysis. *Sports Med* 47: 2341–2361, 2017.
 355. Lally F, Crome P. Understanding frailty. *Postgrad Med J* 83: 16–20, 2007.
 356. Lampe R, Grassl S, Mitternacht J, Gerdesmeyer L, Gradinger R. MRT-measurements of muscle volumes of the lower extremities of youths with spastic hemiplegia caused by cerebral palsy. *Brain Dev* 28: 500–506, 2006.
 357. LaPlante MP, Hendershot GE, Moss AJ. Assistive technology devices and home accessibility features: Prevalence, payment, need, and trends. *Adv Data* 16: 1–11, 1992.
 358. Larsson L. Physical training effects on muscle morphology in sedentary males at different ages. *Med Sci Sports Exerc* 14: 203–206, 1982.
 359. Latham NK, Anderson CS, Lee A, et al. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: The frailty interventions trial in elderly subjects (FITNESS). *J Am Geriatr Soc* 51: 291–299, 2003.
 360. Laughlin GA, Barrett-Connor E. Sexual dimorphism in the influence of advanced aging on adrenal hormone levels: The Rancho Bernardo study. *J Clin Endocrinol Metab* 85: 3561–3568, 2000.
 361. Lavalley ME, Balam T. An overview of strength training injuries: Acute and chronic. *Curr Sports Med Rep* 9: 307–313, 2010.
 362. Law M, Wald N, Morris J. Lowering blood pressure to prevent myocardial infarction and stroke: A new preventive strategy. *Health Technol Assess* 7: 1–94, 2003.
 363. Lazowski DA, Ecclestone NA, Myers AM, et al. A randomized outcome evaluation of group exercise programs in long-term care institutions. *J Gerontol Ser A Biol Sci Med Sci* 54: M621–M8, 1999.
 364. LeBrasseur NK, Walsh K, Arany Z. Metabolic benefits of resistance training and fast glycolytic skeletal muscle. *Am J Physiol Endocrinol Metab* 300: E3–E10, 2011.
 365. Lemmer JT, Hurlbut DE, Martel GF, et al. Age and gender responses to strength training and detraining. *Med Sci Sports Exerc* 32: 1505–1512, 2000.
 366. Lemmer JT, Ivey FM, Ryan AS, et al. Effect of strength training on resting metabolic rate and physical activity: Age and gender comparisons. *Med Sci Sports Exerc* 33: 532–541, 2001.
 367. Leong DP, Teo KK, Rangarajan S, et al. Prognostic value of grip strength: Findings from the prospective Urban Rural epidemiology (PURE) study. *Lancet* 386: 266–273, 2015.
 368. Li WC, Chen YC, Yang RS, Tsauo JY. Effects of exercise programmes on quality of life in osteoporotic and osteopenic postmenopausal women: A systematic review and meta-analysis. *Clin Rehabil* 23: 888–896, 2009.
 369. Liao CD, Tsauo JY, Lin LF, et al. Effects of elastic resistance exercise on body composition and physical capacity in older women with sarcopenic obesity: A CONSORT-compliant prospective randomized controlled trial. *Medicine (Baltimore)* 96: e7115, 2017.
 370. Lim PS, Cheng YM, Wei YH. Large-scale mitochondrial DNA deletions in skeletal muscle of patients with end-stage renal disease. *Free Radic Biol Med* 29: 454–463, 2000.
 371. Lim S, Kim JH, Yoon JW, et al. Sarcopenic obesity: Prevalence and association with metabolic syndrome in the Korean longitudinal study on health and aging (KLoSHA). *Diabetes Care* 33: 1652–1654, 2010.
 372. Lind AR, McNicol GW. Circulatory responses to sustained hand-grip contractions performed during other exercise, both rhythmic and static. *J Physiol* 192: 595–607, 1967.
 373. Lindle RS, Metter EJ, Lynch NA, et al. Age and gender comparisons of muscle strength in 654 women and men aged 20–93 yr. *J Appl Physiol* (1985) 83: 1581–1587, 1997.
 374. Littbrand H, Lundin-Olsson L, Gustafson Y, Rosendahl E. The effect of a high-intensity functional exercise program on activities of daily living: A randomized controlled trial in residential care facilities. *J Am Geriatr Soc* 57: 1741–1749, 2009.
 375. Liu-Ambrose T, Donaldson MG. Exercise and cognition in older adults: Is there a role for resistance training programmes? *Br J Sports Med* 43: 25–27, 2009.
 376. Liu-Ambrose T, Donaldson MG, Ahamed Y, et al. Otago home-based strength and balance retraining improves executive functioning in older fallers: A randomized controlled trial. *J Am Geriatr Soc* 56: 1821–1830, 2008.
 377. Liu-Ambrose TY, Khan KM, Eng JJ, et al. Both resistance and agility training reduce back pain and improve health-related quality of life in older women with low bone mass. *Osteoporos Int* 16: 1321–1329, 2005.
 378. Liu C, Shiroo DM, Jones LY, Clark DO. Systematic review of functional training on muscle strength, physical functioning, and activities of daily living in older adults. *Eur Rev Aging Phys Activity* 11: 95–106, 2014.
 379. Liu CJ, Chang WP, Araujo de Carvalho I, et al. Effects of physical exercise in older adults with reduced physical capacity: meta-analysis of resistance exercise and multimodal exercise. *Int J Rehabil Res* 40: 303–314, 2017.
 380. Liu CJ, Latham N. Can progressive resistance strength training reduce physical disability in older adults? A meta-analysis study. *Disabil Rehabil* 33: 87–97, 2011.
 381. Liu CK, Fielding RA. Exercise as an intervention for frailty. *Clin Geriatr Med* 27: 101–110, 2011.
 382. Liu F, Chang HC. Physiological links of circadian clock and biological clock of aging. *Protein Cell* 8: 477–488, 2017.
 383. Lohne-Seiler H, Torstveit MK, Anderssen SA. Traditional versus functional strength training: Effects on muscle strength and power in the elderly. *J Aging Phys Act* 21: 51–70, 2013.
 384. Lopez P, Pinto RS, Radaelli R, et al. Benefits of resistance training in physically frail elderly: A systematic review. *Aging Clin Exp Res* 30: 889–899, 2018.
 385. Lord SR, Castell S, Corcoran J, et al. The effect of group exercise on physical functioning and falls in frail older people living in retirement villages: A randomized, controlled trial. *J Am Geriatr Soc* 51: 1685–1692, 2003.
 386. Lourenzi FM, Jones A, Pereira DF, et al. Effectiveness of an overall progressive resistance strength program for improving the functional capacity of patients with rheumatoid arthritis: A randomized controlled trial. *Clin Rehabil* 31: 1482–1491, 2017.
 387. Lovell DI, Cuneo R, Gass GC. The effect of strength training and short-term detraining on maximum force and the rate of force development of older men. *Eur J Appl Physiol* 109: 429–435, 2010.
 388. Macaluso A, De Vito G. Muscle strength, power and adaptations to resistance training in older people. *Eur J Appl Physiol* 91: 450–472, 2004.
 389. Macaluso A, De Vito G, Felici F, Nimmo MA. Electromyogram changes during sustained contraction after resistance training in women in their 3rd and 8th decades. *Eur J Appl Physiol* 82: 418–424, 2000.
 390. Macaluso A, Nimmo MA, Foster JE, Cockburn M, McMillan NC, De Vito G. Contractile muscle volume and agonist-antagonist coactivation account for differences in torque between young and older women. *Muscle Nerve* 25: 858–863, 2002.

391. Maganaris CN, Narici MV, Reeves ND. In vivo human tendon mechanical properties: Effect of resistance training in old age. *J Musculoskelet Neuronal Interact* 4: 204–208, 2004.
392. Magni NE, McNair PJ, Rice DA. The effects of resistance training on muscle strength, joint pain, and hand function in individuals with hand osteoarthritis: A systematic review and meta-analysis. *Arthritis Res Ther* 19: 131, 2017.
393. Makizako H, Doi T, Shimada H, et al. Does a multicomponent exercise program improve dual-task performance in amnesic mild cognitive impairment? A randomized controlled trial. *Aging Clin Exp Res* 24: 640–646, 2012.
394. Manini T, Marko M, VanArman T, et al. Efficacy of resistance and task-specific exercise in older adults who modify tasks of everyday life. *J Gerontol A Biol Sci Med Sci* 62: 616–623, 2007.
395. Manini TM, Clark BC. Dynapenia and aging: An update. *J Gerontol A Biol Sci Med Sci* 67: 28–40, 2012.
396. Manini TM, Clark BC, Nalls MA, Goodpaster BH, Ploutz-Snyder LL, Harris TB. Reduced physical activity increases intermuscular adipose tissue in healthy young adults. *Am J Clin Nutr* 85: 377–384, 2007.
397. Marques EA, Mota J, Carvalho J. Exercise effects on bone mineral density in older adults: A meta-analysis of randomized controlled trials. *Age (Dordr)* 34: 1493–1515, 2012.
398. Marsh AP, Miller ME, Rejeski WJ, Hutton SL, Kritchevsky SB. Lower extremity muscle function after strength or power training in older adults. *J Aging Phys Act* 17: 416–443, 2009.
399. Martel GF, Roth SM, Ivey FM, et al. Age and sex affect human muscle fibre adaptations to heavy-resistance strength training. *Exp Physiol* 91: 457–464, 2006.
400. Martínez-Veilla N, Casas-Herrero A, Zambom-Ferraresi F, et al. Effect of exercise intervention on functional decline in very elderly patients during acute hospitalization: A randomized clinical trial. *JAMA Intern Med* 2018. Epub ahead of print.
401. Matheny RW Jr, Nindl BC, Adamo ML. Minireview: Mechano-growth factor: A putative product of IGF-I gene expression involved in tissue repair and regeneration. *Endocrinology* 151: 865–875, 2010.
402. Matthews CE, Chen KY, Freedson PS, et al. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol* 167: 875–881, 2008.
403. Mavros Y, Gates N, Wilson GC, et al. Mediation of cognitive function improvements by strength gains after resistance training in older adults with mild cognitive impairment: Outcomes of the study of mental and resistance training. *J Am Geriatr Soc* 65: 550–559, 2017.
404. McCartney N. Acute responses to resistance training and safety. *Med Sci Sports Exerc* 31: 31–37, 1999.
405. McCartney N, McKelvie RS, Martin J, Sale DG, MacDougall JD. Weight-training-induced attenuation of the circulatory response of older males to weight lifting. *J Appl Physiol* (1985) 74: 1056–1060, 1993.
406. McGrath RP, Kraemer WJ, Vincent BM, Hall OT, Peterson MD. Muscle strength is protective against osteoporosis in an ethnically diverse sample of adults. *J Strength Cond Res* 31: 2586–2589, 2017.
407. McGrath RP, Ottenbacher KJ, Vincent BM, Kraemer WJ, Peterson MD. Muscle weakness and functional limitations in an ethnically diverse sample of older adults. *Ethn Health* 26: 1–12, 2017.
408. McLafferty CL Jr, Wetzstein CJ, Hunter GR. Resistance training is associated with improved mood in healthy older adults. *Percept Mot Skills* 98: 947–957, 2004.
409. McLean RR, Shardell MD, Alley DE, et al. Criteria for clinically relevant weakness and low lean mass and their longitudinal association with incident mobility impairment and mortality: The foundation for the National Institutes of Health (FNIH) sarcopenia project. *J Gerontol A Biol Sci Med Sci* 69: 576–583, 2014.
410. McMurdo ME, Johnstone R. A randomized controlled trial of a home exercise programme for elderly people with poor mobility. *Age Ageing* 24: 425–428, 1995.
411. McMurdo ME, Rennie LM. Improvements in quadriceps strength with regular seated exercise in the institutionalized elderly. *Arch Phys Med Rehabil* 75: 600–603, 1994.
412. McNeil CJ, Vandervoort AA, Rice CL. Peripheral impairments cause a progressive age-related loss of strength and velocity-dependent power in the dorsiflexors. *J Appl Physiol* (1985) 102: 1962–1968, 2007.
413. Melton LJ III, Khosla S, Crowson CS, O'Connor MK, O'Fallon WM, Riggs BL. Epidemiology of sarcopenia. *J Am Geriatr Soc* 48: 625–630, 2000.
414. Metter EJ, Conwit R, Tobin J, Fozard JL. Age-associated loss of power and strength in the upper extremities in women and men. *J Gerontol A Biol Sci Med Sci* 52: B267–B276, 1997.
415. Meuleman JR, Brechue WF, Kubilis PS, Lowenthal DT. Exercise training in the debilitated aged: Strength and functional outcomes. *Arch Phys Med Rehabil* 81: 312–318, 2000.
416. Miszko TA, Cress ME, Slade JM, et al. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 58: 171–175, 2003.
417. Mitch WE. Malnutrition is an unusual cause of decreased muscle mass in chronic kidney disease. *J Ren Nutr* 17: 66–69, 2007.
418. Mitchell JH, Payne FC, Saltin B, Schibye B. The role of muscle mass in the cardiovascular response to static contractions. *J Physiol* 309: 45–54, 1980.
419. Moineddin I, Leehey DJ. A comparison of aerobic exercise and resistance training in patients with and without chronic kidney disease. *Adv Chronic Kidney Dis* 15: 83–96, 2008.
420. Montero-Odasso M, Muir SW, Speechley M. Dual-task complexity affects gait in people with mild cognitive impairment: The interplay between gait variability, dual tasking, and risk of falls. *Arch Phys Med Rehabil* 93: 293–299, 2012.
421. Moore AZ, Caturegli G, Metter EJ, et al. Difference in muscle quality over the adult life span and biological correlates in the Baltimore Longitudinal Study of Aging. *J Am Geriatr Soc* 62: 230–236, 2014.
422. Moraes-Silva IC, Mostarda CT, Silva-Filho AC, Irigoyen MC. Hypertension and exercise training: Evidence from clinical studies. *Adv Exp Med Biol* 1000: 65–84, 2017.
423. Moreau NG, Li L, Geaghan JP, Damiano DL. Fatigue resistance during a voluntary performance task is associated with lower levels of mobility in cerebral palsy. *Arch Phys Med Rehabil* 89: 2011–2016, 2008.
424. Morie M, Reid KF, Miciek R, et al. Habitual physical activity levels are associated with performance in measures of physical function and mobility in older men. *J Am Geriatr Soc* 58: 1727–1733, 2010.
425. Moritani T, deVries HA. Potential for gross muscle hypertrophy in older men. *J Gerontol* 35: 672–682, 1980.
426. Morse CI, Thom JM, Davis MG, et al. Reduced plantarflexor specific torque in the elderly is associated with a lower activation capacity. *Eur J Appl Physiol* 92: 219–226, 2004.
427. Morse CI, Thom JM, Mian OS, Birch KM, Narici MV. Gastrocnemius specific force is increased in elderly males following a 12-month physical training programme. *Eur J Appl Physiol* 100: 563–570, 2007.
428. Morse CI, Thom JM, Mian OS, et al. Muscle strength, volume and activation following 12-month resistance training in 70-year-old males. *Eur J Appl Physiol* 95: 197–204, 2005.
429. Morse CI, Thom JM, Reeves ND, Birch KM, Narici MV. In vivo physiological cross-sectional area and specific force are reduced in the gastrocnemius of elderly men. *J Appl Physiol* (1985) 99: 1050–1055, 2005.
430. Morton RW, Sato K, Gallagher MPB, et al. Muscle androgen receptor content but not systemic hormones is associated with resistance training-induced skeletal muscle hypertrophy in healthy, young men. *Front Physiol* 9: 1373, 2018.
431. Motl RW, McAuley E. Physical activity, disability, and quality of life in older adults. *Phys Med Rehabil Clin N Am* 21: 299–308, 2010.
432. Mozaffarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statistics-2015 update a report from the American heart association. *Circulation* 131: E29–E322, 2015.
433. Muhlberg W, Sieber C. Sarcopenia and frailty in geriatric patients: Implications for training and prevention. *Z Gerontol Geriatr* 37: 2–8, 2004.
434. Nader GA. Concurrent strength and endurance training: From molecules to man. *Med Sci Sports Exerc* 38: 1965–1970, 2006.
435. Nakamura Y, Tanaka K, Yabushita N, Sakai T, Shigematsu R. Effects of exercise frequency on functional fitness in older adult women. *Arch Gerontol Geriatr* 44: 163–173, 2007.
436. Narici MV, Maganaris CN, Reeves ND, Capodaglio P. Effect of aging on human muscle architecture. *J Appl Physiol* (1985) 95: 2229–2234, 2003.
437. Narici MV, Reeves ND, Morse CI, Maganaris CN. Muscular adaptations to resistance exercise in the elderly. *J Musculoskelet Neuronal Interact* 4: 161–164, 2004.
438. Nelson ME, Fiatarone MA, Morganti CM, et al. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA* 272: 1909–1914, 1994.
439. Nelson RM, Soderberg GL, Urbscheit NL. Alteration of motor-unit discharge characteristics in aged humans. *Phys Ther* 64: 29–34, 1984.
440. Neves SdaC Jr, Lima RM, Simoes HG, et al. Resistance exercise sessions do not provoke acute immunosuppression in older women. *J Strength Cond Res* 23: 259–265, 2009.
441. Newton RU, Hakkinen K, Hakkinen A, McCormick M, Volek J, Kraemer WJ. Mixed-methods resistance training increases power and strength of young and older men. *Med Sci Sports Exerc* 34: 1367–1375, 2002.
442. Nguyen C, Lefevre-Colau MM, Poiraudou S, Rannou F. Rehabilitation (exercise and strength training) and osteoarthritis: A critical narrative review. *Ann Phys Rehabil Med* 59: 190–195, 2016.
443. Nicklas BJ, Brinkley TE. Exercise training as a treatment for chronic inflammation in the elderly. *Exerc Sport Sci Rev* 37: 165–170, 2009.
444. National Institute of Diabetes and Digestive and Kidney Diseases. Kidney Disease Statistics for the United States. Atlanta, GA: Centers for Disease Control and Prevention, 2016.
445. Nielson CM, Srikanth P, Orwoll ES. Obesity and fracture in men and women: An epidemiologic perspective. *J Bone Miner Res* 27: 1–10, 2012.
446. Nindl BC. Insulin-like growth factor-1, physical activity, and control of cellular anabolism. *Med Sci Sports Exerc* 42: 35–38, 2010.
447. Nindl BC, Kraemer WJ, Marx JO, Tuckow AP, Hymer WC. Growth hormone molecular heterogeneity and exercise. *Exerc Sport Sci Rev* 31: 161–166, 2003.
448. Nindl BC, Pierce JR. Insulin-like growth factor 1 as a biomarker of health, fitness, and training status. *Med Sci Sports Exerc* 42: 39–49, 2010.
449. NSCA. Essentials of Strength Training and Conditioning. Hum Kinetics, 2016. pp. 462.
450. O'Connor KG, Tobin JD, Harman SM, et al. Serum levels of insulin-like growth factor-I are related to age and not to body composition in healthy women and men. *J Gerontol A Biol Sci Med Sci* 53: M176–M182, 1998.

451. Palmer TB, Thiele RM, Conchola EC, Smith DB, Thompson BJ. A preliminary study of the utilization of maximal and rapid strength characteristics to identify chair-rise performance abilities in very old adults. *J Geriatr Phys Ther* 39: 102–109, 2016.
452. Palmer TB, Thiele RM, Thompson BJ. Age-related differences in maximal and rapid torque characteristics of the hip extensors and dynamic postural balance in healthy, young and old females. *J Strength Cond Res* 31: 480–488, 2017.
453. Palmer TB, Thiele RM, Williams KB, et al. The identification of fall history using maximal and rapid isometric torque characteristics of the hip extensors in healthy, recreationally active elderly females: A preliminary investigation. *Aging Clin Exp Res* 27: 431–438, 2015.
454. Pantoni LB, Franke WD, Bleil DA, Baier SM, King DS. Effects of resistance training on cardiovascular responses to lower body negative pressure in the elderly. *Clin Physiol* 21: 605–611, 2001.
455. Papa EV, Dong X, Hassan M. Resistance training for activity limitations in older adults with skeletal muscle function deficits: A systematic review. *Clin Interv Aging* 12: 955–961, 2017.
456. Papa EV, Dong X, Hassan M. Skeletal muscle function deficits in the elderly: Current perspectives on resistance training. *J Nat Sci* 3: pii: e272, 2017.
457. Park J, Mendy A, Vieira ER. Various types of arthritis in the United States: Prevalence and age-related trends from 1999 to 2014. *Am J Public Health* 108: 256–258, 2018.
458. Pasco JA, Gould H, Brennan SL, Nicholson GC, Kotowicz MA. Musculoskeletal deterioration in men accompanies increases in body fat. *Obesity (Silver Spring)* 22: 863–867, 2014.
459. Pearson SJ, Young A, Macaluso A, et al. Muscle function in elite master weightlifters. *Med Sci Sports Exerc* 34: i199–i206, 2002.
460. Penninx BW, Kritchevsky SB, Newman AB, et al. Inflammatory markers and incident mobility limitation in the elderly. *J Am Geriatr Soc* 52: 1105–1113, 2004.
461. Penninx BW, Messier SP, Rejeski WJ, et al. Physical exercise and the prevention of disability in activities of daily living in older persons with osteoarthritis. *Arch Intern Med* 161: 2309–2316, 2001.
462. Peterson M, Duchowny K, Meng Q, et al. Low normalized grip strength is a biomarker for cardiometabolic disease and physical disabilities among U.S. and Chinese adults. *J Gerontol A Biol Sci Med Sci* 72: 1525–1531, 2017.
463. Peterson MD, Belakovskiy A, McGrath R, Yarrow JF. Testosterone deficiency, weakness, and multimorbidity in men. *Sci Rep* 8: 5897, 2018.
464. Peterson MD, Mahmoudi E. Healthcare utilization associated with obesity and physical disabilities. *Am J Prev Med* 48: 426–435, 2015.
465. Peterson MD, Rhea MR, Sen A, Gordon PM. Resistance exercise for muscular strength in older adults: A meta-analysis. *Ageing Res Rev* 9: 226–237, 2010.
466. Peterson MD, Ryan JM, Hurvitz EA, Mahmoudi E. Chronic conditions in adults with cerebral palsy. *JAMA* 314: 2303–2305, 2015.
467. Peterson MD, Saltarelli WA, Visich PS, Gordon PM. Strength capacity and cardiometabolic risk clustering in adolescents. *Pediatrics* 133: e896–903, 2014.
468. Peterson MD, Sen A, Gordon PM. Influence of resistance exercise on lean body mass in aging adults: A meta-analysis. *Med Sci Sports Exerc* 43: 249–258, 2011.
469. Peterson MD, Zhang P, Choksi P, Markides KS, Al Snih S. Muscle weakness thresholds for prediction of diabetes in adults. *Sports Med* 46: 619–628, 2016.
470. Peterson MD, Zhang P, Duchowny KA, et al. Declines in strength and mortality risk among older Mexican Americans: Joint modeling of survival and longitudinal data. *J Gerontol A Biol Sci Med Sci* 71: 1646–1652, 2016.
471. Peterson MD, Zhang P, Haapala HJ, Wang SC, Hurvitz EA. Greater adipose tissue distribution and diminished spinal musculoskeletal density in adults with cerebral palsy. *Arch Phys Med Rehabil* 96: 1828–1833, 2015.
472. Petrella JK, Kim JS, Cross JM, Kosek DJ, Bamman MM. Efficacy of myonuclear addition may explain differential myofiber growth among resistance-trained young and older men and women. *Am J Physiol Endocrinol Metab* 291: E937–E946, 2006.
473. Phillips SK, Bruce SA, Newton D, Woledge RC. The weakness of old age is not due to failure of muscle activation. *J Gerontol* 47: M45–M49, 1992.
474. Phillips SM. Resistance exercise: Good for more than just grandma and grandpa's muscles. *Appl Physiol Nutr Metab* 32: 1198–1205, 2007.
475. Piepoli MF, Conraads V, Corra U, et al. Exercise training in heart failure: From theory to practice. A consensus document of the heart failure association and the European association for cardiovascular prevention and rehabilitation. *Eur J Heart Fail* 13: 347–357, 2011.
476. Pina IL, Apstein CS, Balady GJ, et al. Exercise and heart failure: A statement from the American heart association Committee on exercise, rehabilitation, and prevention. *Circulation* 107: 1210–1225, 2003.
477. Pinto RS, Correa CS, Radaelli R, et al. Short-term strength training improves muscle quality and functional capacity of elderly women. *Age (Dordr)* 36: 365–372, 2014.
478. Pitkälä K, Savikko N, Poysti M, Strandberg T, Laakkonen ML. Efficacy of physical exercise intervention on mobility and physical functioning in older people with dementia: A systematic review. *Exp Gerontol* 48: 85–93, 2013.
479. Plassman BL, Langa KM, Fisher GG, et al. Prevalence of dementia in the United States: The aging, demographics, and memory study. *Neuro-epidemiology* 29: 125–132, 2007.
480. Pollock ML, Franklin BA, Balady GJ, et al. AHA/Science Advisory. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription: An advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association; Position paper endorsed by the American College of Sports Medicine. *Circulation* 101: 828–833, 2000.
481. Price SR, Gooch JL, Donaldson SK, Roberts-Wilson TK. Muscle atrophy in chronic kidney disease results from abnormalities in insulin signaling. *J Ren Nutr* 20(5 Suppl): S24–S28, 2010.
482. Pu CT, Johnson MT, Forman DE, et al. Randomized trial of progressive resistance training to counteract the myopathy of chronic heart failure. *J Appl Physiol* (1985) 90: 2341–2350, 2001.
483. Pyka G, Lindenberger E, Charette S, Marcus R. Muscle strength and fiber adaptations to a year-long resistance training program in elderly men and women. *J Gerontol* 49: M22–M27, 1994.
484. Pyka G, Taaffe DR, Marcus R. Effect of a sustained program of resistance training on the acute growth hormone response to resistance exercise in older adults. *Horm Metab Res* 26: 330–333, 1994.
485. Qureshi AR, Alvestrand A, Danielsson A, et al. Factors predicting malnutrition in hemodialysis patients: A cross-sectional study. *Kidney Int* 53: 773–782, 1998.
486. Radaelli R, Botton CE, Wilhelm EN, et al. Time course of low- and high-volume strength training on neuromuscular adaptations and muscle quality in older women. *Age (Dordr)* 36: 881–892, 2014.
487. Radaelli R, Brusco CM, Lopez P, et al. Higher muscle power training volume is not determinant for the magnitude of neuromuscular improvements in elderly women. *Exp Gerontol* 110: 15–22, 2018.
488. Ramirez-Campillo R, Castillo A, de la Fuente CI, et al. High-speed resistance training is more effective than low-speed resistance training to increase functional capacity and muscle performance in older women. *Exp Gerontol* 58: 51–57, 2014.
489. Ramsbottom R, Ambler A, Potter J, Jordan B, Nevill A, Williams C. The effect of 6 months training on leg power, balance, and functional mobility of independently living adults over 70 years old. *J Aging Phys Act* 12: 497–510, 2004.
490. Rantanen T, Guralnik JM, Ferrucci L, Leveille S, Fried LP. Compairments: Strength and balance as predictors of severe walking disability. *J Gerontol A Biol Sci Med Sci* 54: M172–M176, 1999.
491. Raso V, Benard G, DA Silva Duarte AJ, Natale VM. Effect of resistance training on immunological parameters of healthy elderly women. *Med Sci Sports Exerc* 39: 2152–2159, 2007.
492. Raue U, Slivka D, Minchev K, Trappe S. Improvements in whole muscle and myocellular function are limited with high-intensity resistance training in octogenarian women. *J Appl Physiol* (1985) 106: 1611–1617, 2009.
493. Raymond MJ, Bramley-Tzerofos RE, Jeffs KJ, Winter A, Holland AE. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. *Arch Phys Med Rehabil* 94: 1458–1472, 2013.
494. Reed RL, Pearlmutter L, Yochum K, Meredith KE, Mooradian AD. The relationship between muscle mass and muscle strength in the elderly. *J Am Geriatr Soc* 39: 555–561, 1991.
495. Reed T, Fabsitz RR, Selby JV, Carmelli D. Genetic influences and grip strength norms in the NHLBI twin study males aged 59–69. *Ann Hum Biol* 18: 425–432, 1991.
496. Reeves ND, Narici MV, Maganaris CN. Effect of resistance training on skeletal muscle-specific force in elderly humans. *J Appl Physiol* (1985) 96: 885–892, 2004.
497. Reeves ND, Narici MV, Maganaris CN. In vivo human muscle structure and function: Adaptations to resistance training in old age. *Exp Physiol* 89: 675–689, 2004.
498. Reeves ND, Narici MV, Maganaris CN. Musculoskeletal adaptations to resistance training in old age. *Man Ther* 11: 192–196, 2006.
499. Reid KF, Callahan DM, Carabello RJ, et al. Lower extremity power training in elderly subjects with mobility limitations: A randomized controlled trial. *Aging Clin Exp Res* 20: 337–343, 2008.
500. Reid KF, Fielding RA. Skeletal muscle power: A critical determinant of physical functioning in older adults. *Exerc Sport Sci Rev* 40: 4–12, 2012.
501. Reid KF, Martin KI, Doros G, et al. Comparative effects of light or heavy resistance power training for improving lower extremity power and physical performance in mobility-limited older adults. *J Gerontol A Biol Sci Med Sci* 70: 374–380, 2015.
502. Reynolds TH IV, Supiano MA, Dengel DR. Regional differences in glucose clearance: Effects of insulin and resistance training on arm and leg glucose clearance in older hypertensive individuals. *J Appl Physiol* (1985) 102: 985–991, 2007.
503. Rhodes EC, Martin AD, Taunton JE, et al. Effects of one year of resistance training on the relation between muscular strength and bone density in elderly women. *Br J Sports Med* 34: 18–22, 2000.
504. Ribeiro AS, Schoenfeld BJ, Fleck SJ, et al. Effects of traditional and pyramidal resistance training systems on muscular strength, muscle mass, and hormonal responses in older women: A randomized crossover trial. *J Strength Cond Res* 31: 1888–1896, 2017.
505. Ribeiro F, Costa R, Mesquita-Bastos J. Exercise training in the management of patients with resistant hypertension. *World J Cardiol* 7: 47–51, 2015.
506. Riebe D, Franklin BA, Thompson PD, et al. Updating ACSM's recommendations for exercise preparticipation health screening. *Med Sci Sports Exerc* 47: 2473–2479, 2015.
507. Robergs RA, Ghiasvand F, Parker D. Biochemistry of exercise-induced metabolic acidosis. *Am J Physiol Regul Integr Comp Physiol* 287: R502–R516, 2004.

508. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci* 62: 722–727, 2007.
509. Rodríguez-Manas L, Fried LP. Frailty in the clinical scenario. *Lancet* 385: e7–e9, 2015.
510. Roman WJ, Fleckenstein J, Stray-Gundersen J, et al. Adaptations in the elbow flexors of elderly males after heavy-resistance training. *J Appl Physiol* (1985) 74: 750–754, 1993.
511. Rooyackers OE, Adey DB, Ades PA, Nair KS. Effect of age on in vivo rates of mitochondrial protein synthesis in human skeletal muscle. *Proc Natl Acad Sci U S A* 93: 15364–15369, 1996.
512. Rose J, McGill KC. Neuromuscular activation and motor-unit firing characteristics in cerebral palsy. *Dev Med Child Neurol* 47: 329–336, 2005.
513. Rose SJ, Rothstein JM. Muscle mutability. Part I. General concepts and adaptations to altered patterns of use. *Phys Ther* 62: 1773–1787, 1982.
514. Rosendahl E, Lindelo f N, Littbrand H, et al. High-intensity functional exercise program and protein-enriched energy supplement for older persons dependent in activities of daily living: A randomised controlled trial. *Aust J Physiother* 52: 105–113, 2006.
515. Rossi A, Zoico E, Goodpaster BH, et al. Quantification of intermuscular adipose tissue in the Erector spinae muscle by MRI: Agreement with histological evaluation. *Obesity (Silver Spring)* 18: 2379–2384, 2010.
516. Roth SM, Ivey FM, Martel GF, et al. Muscle size responses to strength training in young and older men and women. *J Am Geriatr Soc* 49: 1428–1433, 2001.
517. Rothstein JM, Rose SJ. Muscle mutability. Part 2. Adaptation to drugs, metabolic factors, and aging. *Phys Ther* 62: 1788–1798, 1982.
518. Roubenoff R. Sarcopenia: A major modifiable cause of frailty in the elderly. *J Nutr Health Aging* 4: 140–142, 2000.
519. Roubenoff R, Hughes VA. Sarcopenia: Current concepts. *J Gerontol A Biol Sci Med Sci* 55: M716–M724, 2000.
520. Rubin MR, Kraemer WJ, Kraemer RR, et al. Responses of growth hormone aggregates to different intermittent exercise intensities. *Eur J Appl Physiol* 89: 166–170, 2003.
521. Ruiz J, Sui X, Lobelo F, et al. Association between muscular strength and mortality in men: Prospective cohort study. *BMJ* 337: a439, 2008.
522. Ryu M, Jo J, Lee Y, Chung YS, Kim KM, Baek WC. Association of physical activity with sarcopenia and sarcopenic obesity in community-dwelling older adults: The fourth Korea National health and nutrition examination survey. *Age and Ageing* 42: 734–740, 2013.
523. Sambrook P, Cooper C. Osteoporosis. *Lancet* 367: 2010–2018, 2006.
524. Sanchez-Medina L, Gonzalez-Badillo JJ. Velocity loss as an indicator of neuromuscular fatigue during resistance training. *Med Sci Sports Exerc* 43: 1725–1734, 2011.
525. Sardeli AV, Tomeleri CM, Cyrino ES, et al. Effect of resistance training on inflammatory markers of older adults: A meta-analysis. *Exp Gerontol* 111: 188–196, 2018.
526. Sayer AA, Kirkwood TB. Grip strength and mortality: A biomarker of ageing? *Lancet* 386: 226–227, 2015.
527. Sayers SP. High velocity power training in older adults. *Curr Aging Sci* 1: 62–67, 2008.
528. Sayers SP, Gibson K. A comparison of high-speed power training and traditional slow-speed resistance training in older men and women. *J Strength Cond Res* 24: 3369–3380, 2010.
529. Scaglioni G, Ferri A, Minetti AE, et al. Plantar flexor activation capacity and H reflex in older adults: Adaptations to strength training. *J Appl Physiol* (1985) 92: 2292–2302, 2002.
530. Scher LM, Ferrioli E, Moriguti JC, Scher R, Lima NK. The effect of different volumes of acute resistance exercise on elderly individuals with treated hypertension. *J Strength Cond Res* 25: 1016–1023, 2011.
531. Schoenfeld BJ. Postexercise hypertrophic adaptations: A reexamination of the hormone hypothesis and its applicability to resistance training program design. *J Strength Cond Res* 27: 1720–1730, 2013.
532. Schraeger MA, Metter EJ, Simonsick E, et al. Sarcopenic obesity and inflammation in the InCHIANTI study. *J Appl Physiol* (1985) 102: 919–925, 2007.
533. Seguin R, Nelson ME. The benefits of strength training for older adults. *Am J Prev Med* 25(3 Suppl 2): 141–149, 2003.
534. Seidler RD, Bernard JA, Burutolu TB, et al. Motor control and aging: Links to age-related brain structural, functional, and biochemical effects. *Neurosci Biobehav Rev* 34: 721–733, 2010.
535. Senechal M, McGavock JM, Church TS, et al. Cut points of muscle strength associated with metabolic syndrome in men. *Med Sci Sports Exerc* 46: 1475–1481, 2014.
536. Serra-Rexach JA, Bustamante-Ara N, Hierro Villaran M, et al. Short-term, light-to moderate-intensity exercise training improves leg muscle strength in the oldest old: A randomized controlled trial. *J Am Geriatr Soc* 59: 594–602, 2011.
537. Seynnes O, Fiatarone Singh MA, Hue O, et al. Physiological and functional responses to low-moderate versus high-intensity progressive resistance training in frail elders. *J Gerontol A Biol Sci Med Sci* 59: 503–509, 2004.
538. Shafiqe G, Keshitkar A, Soltani A, et al. Prevalence of sarcopenia in the world: A systematic review and meta-analysis of general population studies. *J Diabetes Metab Disord* 16: 21, 2017.
539. Shah K, Armamento-Villareal R, Parimi N, et al. Exercise training in obese older adults prevents increase in bone turnover and attenuates decrease in hip bone mineral density induced by weight loss despite decline in bone-active hormones. *J Bone Miner Res* 26: 2851–2859, 2011.
540. Sharman MJ, Newton RU, Triplett-McBride T, et al. Changes in myosin heavy chain composition with heavy resistance training in 60- to 75- year-old men and women. *Eur J Appl Physiol* 84: 127–132, 2001.
541. Shaw BS, Shaw I. Effect of resistance training on cardiorespiratory endurance and coronary artery disease risk. *Cardiovasc J S Afr* 16: 256–259, 2005.
542. Shaw BS, Shaw I, Brown GA. Resistance exercise is medicine: Strength training in health promotion and rehabilitation. *Int J Ther Rehabil* 22: 385–389, 2015.
543. Shaw CS, Clark J, Wagenmakers AJ. The effect of exercise and nutrition on intramuscular fat metabolism and insulin sensitivity. *Annu Rev Nutr* 30: 13–34, 2010.
544. Sheridan KJ. Osteoporosis in adults with cerebral palsy. *Dev Med Child Neurol* 51(Suppl 4): 38–51, 2009.
545. Sherrington C, Michaleff ZA, Fairhall N, et al. Exercise to prevent falls in older adults: An updated systematic review and meta-analysis. *Br J Sports Med* 51: 1750–1758, 2017.
546. Sherrington C, Whitney JC, Lord SR, et al. Effective exercise for the prevention of falls: A systematic review and meta-analysis. *J Am Geriatr Soc* 56: 2234–2243, 2008.
547. Shinohara M. Adaptations in motor unit behavior in elderly adults. *Curr Aging Sci* 4: 200–208, 2011.
548. Shiroma EJ, Cook NR, Manson JE, et al. Strength training and the risk of type 2 diabetes and cardiovascular disease. *Med Sci Sports Exerc* 49: 40–46, 2017.
549. Short KR, Nair KS. Mechanisms of sarcopenia of aging. *J Endocrinol Invest* 22(5 Suppl): 95–105, 1999.
550. Shortland A. Muscle deficits in cerebral palsy and early loss of mobility: Can we learn something from our elders? *Dev Med Child Neurol* 51(- Suppl 4): 59–63, 2009.
551. Sillanpaa E, Hakkinen A, Nyman K, et al. Body composition and fitness during strength and/or endurance training in older men. *Med Sci Sports Exerc* 40: 950–958, 2008.
552. Silva NL, Oliveira RB, Fleck SJ, Leon AC, Farinatti P. Influence of strength training variables on strength gains in adults over 55 years-old: A meta-analysis of dose-response relationships. *J Sci Med Sport* 17: 337–344, 2014.
553. Silva RB, Eslick GD, Duque G. Exercise for falls and fracture prevention in long term care facilities: A systematic review and meta-analysis. *J Am Med Dir Assoc* 14: 685–689.e2, 2013.
554. Simao R, de Salles BF, Figueiredo T, Dias I, Willardson JM. Exercise order in resistance training. *Sports Med* 42: 251–265, 2012.
555. Singh MA. Exercise comes of age: Rationale and recommendations for a geriatric exercise prescription. *J Gerontol A Biol Sci Med Sci* 57: M262–M282, 2002.
556. Singh NA, Stavrinou TM, Scarbek Y, et al. A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults. *J Gerontol A Biol Sci Med Sci* 60: 768–776, 2005.
557. Sipe C, Ritchie D. The significant 7 principles of functional training for mature adults. *IDEA Fitness J* 9: 42–49, 2012.
558. Skelton DA, Greig CA, Davies JM, Young A. Strength, power and related functional ability of healthy people aged 65–89 years. *Age Ageing* 23: 371–377, 1994.
559. Skelton DA, Young A, Greig CA, Malbut KE. Effects of resistance training on strength, power, and selected functional abilities of women aged 75 and older. *J Am Geriatr Soc* 43: 1081–1087, 1995.
560. Slemenda C, Brandt KD, Heilman DK, et al. Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med* 127: 97–104, 1997.
561. Slivka D, Raue U, Hollon C, Minchev K, Trappe S. Single muscle fiber adaptations to resistance training in old (.80 yr) men: Evidence for limited skeletal muscle plasticity. *Am J Physiol Regul Integr Comp Physiol* 295: R273–R280, 2008.
562. Sousa N, Mendes R, Abrantes C, Sampaio J, Oliveira J. Effectiveness of combined exercise training to improve functional fitness in older adults: A randomized controlled trial. *Geriatr Gerontol Int* 14: 892–898, 2014.
563. Sousa N, Mendes R, Monteiro G, Abrantes C. Progressive resistance strength training and the related injuries in older adults: The susceptibility of the shoulder. *Aging Clin Exp Res* 26: 235–240, 2014.
564. Sousa N, Mendes R, Silva A, Oliveira J. Combined exercise is more effective than aerobic exercise in the improvement of fall risk factors: A randomized controlled trial in community-dwelling older men. *Clin Rehabil* 31: 478–486, 2017.
565. Souza AH, Farias MI, Salvatori R, et al. Lifetime, untreated isolated GH deficiency due to a GH-releasing hormone receptor mutation has beneficial consequences on bone status in older individuals, and does not influence their abdominal aorta calcification. *Endocrine* 47: 191–197, 2014.
566. Spiering BA, Kraemer WJ, Anderson JM, et al. Resistance exercise biology: Manipulation of resistance exercise programme variables determines the responses of cellular and molecular signalling pathways. *Sports Med* 38: 527–540, 2008.
567. Spirduso WW, Cronin DL. Exercise dose-response effects on quality of life and independent living in older adults. *Med Sci Sports Exerc* 33(6 Suppl): S598–S608, 2001; discussion S9–10.
568. Spruit MA, Sillen MJH, Groenen MTJ, Wouters EFM, Franssen FME. New normative values for handgrip strength: Results from the UK Bio-bank. *J Am Med Dir Assoc* 14: 775.e5–775.e11, 2013.
569. Stackhouse SK, Stapleton MR, Wagner DA, McClure PW. Voluntary activation of the infraspinatus muscle in nonfatigued and fatigued states. *J Shoulder Elbow Surg* 19: 224–229, 2010.
570. National Center for Health Statistics. Survey Description, National Health Interview Survey, 2015. Hyattsville, Maryland: National Center for Health Statistics, 2016.

571. Steele J, Bruce-Low S, Smith D, Jessop D, Osborne N. A randomized controlled trial of limited range of motion lumbar extension exercise in chronic low back pain. *Spine (Phila Pa 1976)* 38: 1245–1252, 2013.
572. Steffens DC, Fisher GG, Langa KM, Potter GG, Plassman BL. Prevalence of depression among older Americans: The aging, demographics and memory study. *Int Psychogeriatr* 21: 879–888, 2009.
573. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults: A meta-analysis. *Med Sci Sports Exerc* 42: 902–914, 2010.
574. Stenholm S, Harris TB, Rantanen T, et al. Sarcopenic obesity: Definition, cause and consequences. *Curr Opin Clin Nutr Metab Care* 11: 693–700, 2008.
575. Stenroth L, Peltonen J, Cronin NJ, Sipilä S, Finni T. Age-related differences in Achilles tendon properties and triceps surae muscle architecture in vivo. *J Appl Physiol (1985)* 113: 1537–1544, 2012.
576. Stensvold D, Tjønnå AE, Skaug EA, et al. Strength training versus aerobic interval training to modify risk factors of metabolic syndrome. *J Appl Physiol (1985)* 108: 804–810, 2010.
577. Stevens LA, Viswanathan G, Weiner DE. Chronic kidney disease and end-stage renal disease in the elderly population: Current prevalence, future projections, and clinical significance. *Adv Chronic Kidney Dis* 17: 293–301, 2010.
578. Stevenson LW, Massie BM, Francis GS. Optimizing therapy for complex or refractory heart failure: A management algorithm. *Am Heart J* 135(6 Pt 2 Suppl): S293–S309, 1998.
579. Stewart KJ, Bacher AC, Turner KL, et al. Effect of exercise on blood pressure in older persons: A randomized controlled trial. *Arch Intern Med* 165: 756–762, 2005.
580. Straight CR, Lindheimer JB, Brady AO, Dishman RK, Evans EM. Effects of resistance training on lower-extremity muscle power in middle-aged and older adults: A systematic review and meta-analysis of randomized controlled trials. *Sports Med* 46: 353–364, 2016.
581. Strasser B, Keinrad M, Haber P, Schobersberger W. Efficacy of systematic endurance and resistance training on muscle strength and endurance performance in elderly adults—A randomized controlled trial. *Wien Klin Wochenschr* 121: 757–764, 2009.
582. Strasser B, Schobersberger W. Evidence for resistance training as a treatment therapy in obesity. *J Obes* 2011: pii: 482564, 2011.
583. Suetta C, Aagaard P, Rosted A, et al. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *J Appl Physiol (1985)* 97: 1954–1961, 2004.
584. Suetta C, Andersen JL, Dalgaard U, et al. Resistance training induces qualitative changes in muscle morphology, muscle architecture, and muscle function in elderly postoperative patients. *J Appl Physiol (1985)* 105: 180–186, 2008.
585. Suetta C, Hvid LG, Justesen L, et al. Effects of aging on human skeletal muscle after immobilization and retraining. *J Appl Physiol (1985)* 107: 1172–1180, 2009.
586. Sullivan DH, Roberson PK, Smith ES, Price JA, Bopp MM. Effects of muscle strength training and megesterol acetate on strength, muscle mass, and function in frail older people. *J Am Geriatr Soc* 55: 20–28, 2007.
587. Sullivan DH, Wall PT, Bariola JR, Bopp MM, Frost YM. Progressive resistance muscle strength training of hospitalized frail elderly. *Am J Phys Med Rehabil* 80: 503–509, 2001.
588. Suzuki T, Shimada H, Makizako H, et al. Effects of multicomponent exercise on cognitive function in older adults with amnesic mild cognitive impairment: A randomized controlled trial. *BMC Neurol* 12: 128, 2012.
589. Taaffe DR, Duret C, Wheeler S, Marcus R. Once-weekly resistance exercise improves muscle strength and neuromuscular performance in older adults. *J Am Geriatr Soc* 47: 1208–1214, 1999.
590. Taekema DG, Gussekloo J, Maier AB, Westendorp RGJ, de Craen AJM. Handgrip strength as a predictor of functional, psychological and social health. A prospective population-based study among the oldest old. *Age Ageing* 39: 331–337, 2010.
591. Tanasescu M, Leitzmann MF, Rimm EB, et al. Exercise type and intensity in relation to coronary heart disease in men. *JAMA* 288: 1994–2000, 2002.
592. Tapps T, Passmore T, Lindenmeier D, Bishop A. An investigation into the effects of resistance based physical activity participation on depression of older adults in a long-term care facility. *Annu Ther Recreat* 21: 63–72, 2013.
593. Tarazona-Santabalbina FJ, Gomez-Cabrera MC, Perez-Ros P, et al. A multicomponent exercise intervention that reverses frailty and improves cognition, emotion, and social networking in the community-dwelling frail elderly: A randomized clinical trial. *J Am Med Dir Assoc* 17: 426–433, 2016.
594. Tarnopolsky M, Zimmer A, Paikin J, et al. Creatine monohydrate and conjugated linoleic acid improve strength and body composition following resistance exercise in older adults. *PLoS One* 2: e991, 2007.
595. Taylor D, Hale L, Schluter P, et al. Effectiveness of tai chi as a community-based falls prevention intervention: A randomized controlled trial. *J Am Geriatr Soc* 60: 841–848, 2012.
596. Tesch PA, Fernandez-Gonzalo R, Lundberg TR. Clinical applications of iso-inertial, eccentric-overload (YoYo) resistance exercise. *Front Physiol* 8: 241, 2017.
597. Tetsuka K, Hoshi T, Sumiya E, et al. The influence of aging on renal blood flow in human beings. *J Med Ultrason (2001)* 30: 247–251, 2003.
598. Teumer A, Qi Q, Nethander M, et al. Genomewide meta-analysis identifies loci associated with IGF-1 and IGFBP-3 levels with impact on age-related traits. *Aging Cell* 15: 811–824, 2016.
599. Teychenne M, Ball K, Salmon J. Sedentary behavior and depression among adults: A review. *Int J Behav Med* 17: 246–254, 2010.
600. Theou O, Jones GR, Vandervoort AA, Jakobi JM. Daily muscle activity and quiescence in non-frail, pre-frail, and frail older women. *Exp Gerontol* 45: 909–917, 2010.
601. Thomis MA, Beunen GP, Maes HH, et al. Strength training: Importance of genetic factors. *Med Sci Sports Exerc* 30: 724–731, 1998.
602. Thompson BJ, Ryan ED, Herda TJ, Costa PB, Herda AA, Cramer JT. Age-related changes in the rate of muscle activation and rapid force characteristics. *Age (Dordr)* 36: 839–849, 2014.
603. Thompson BJ, Ryan ED, Sobolewski EJ, Conchola EC, Cramer JT. Age related differences in maximal and rapid torque characteristics of the leg extensors and flexors in young, middle-aged and old men. *Exp Gerontol* 48: 277–282, 2013.
604. Tomlinson DJ, Erskine RM, Winwood K, Morse CI, Onambele GL. The impact of obesity on skeletal muscle architecture in untrained young vs. old women. *J Anat* 225: 675–684, 2014.
605. Topp R, Boardley D, Morgan AL, Fahlman M, McNevin N. Exercise and functional tasks among adults who are functionally limited. *West J Nurs Res* 27: 252–270, 2005.
606. Tracy BL, Byrnes WC, Enoka RM. Strength training reduces force fluctuations during anisometric contractions of the quadriceps femoris muscles in old adults. *J Appl Physiol (1985)* 96: 1530–1540, 2004.
607. Tracy BL, Ivey FM, Hurlbut D, et al. Muscle quality. II. Effects of strength training in 65- to 75-year-old men and women. *J Appl Physiol (1985)* 86: 195–201, 1999.
608. Trappe S, Godard M, Gallagher P, et al. Resistance training improves single muscle fiber contractile function in older women. *Am J Physiol Cell Physiol* 281: C398–C406, 2001.
609. Trappe S, Williamson D, Godard M, Porter D, Rowden G, Costill D. Effect of resistance training on single muscle fiber contractile function in older men. *J Appl Physiol (1985)* 89: 143–152, 2000.
610. Treadwell JR, Singh S, Talati R, McPheeters ML, Reston JT. A Framework for “Best Evidence” Approaches in Systematic Reviews. Rockville, MD: Agency for Healthcare Research and Quality (US), 2011. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK56653/>. Accessed June 11, 2018.
611. Treuth MS, Ryan AS, Pratley RE, et al. Effects of strength training on total and regional body composition in older men. *J Appl Physiol (1985)* 77: 614–620, 1994.
612. Trinh A, Wong P, Fahey MC, et al. Musculoskeletal and endocrine health in adults with cerebral palsy: New opportunities for intervention. *J Clin Endocrinol Metab* 101: 1190–1197, 2016.
613. Trouwborst I, Verreijen A, Memelink R, et al. Exercise and nutrition strategies to counteract sarcopenic obesity. *Nutrients* 10: pii: E605, 2018.
614. Tsai CL, Wang CH, Pan CY, Chen FC. The effects of long-term resistance exercise on the relationship between neurocognitive performance and GH, IGF-1, and homocysteine levels in the elderly. *Front Behav Neurosci* 9: 23, 2015.
615. Tschopp M, Sattelmayer MK, Hilfiker R. Is power training or conventional resistance training better for function in elderly persons? A meta-analysis. *Age Ageing* 40: 549–556, 2011.
616. Tsutsumi T, Don BM, Zaichkowsky LD, Delizonna LL. Physical fitness and psychological benefits of strength training in community dwelling older adults. *Appl Hum Sci* 16: 257–266, 1997.
617. Tsutsumi T, Don BM, Zaichkowsky LD, Takenaka K, Oka K, Ohno T. Comparison of high and moderate intensity of strength training on mood and anxiety in older adults. *Percept Mot Skills* 87: 1003–1011, 1998.
618. Tuttle LJ, Sinacore DR, Mueller MJ. Intermuscular adipose tissue is muscle specific and associated with poor functional performance. *J Aging Research* 2012: 172957, 2012.
619. Unhjem R, Lundestad R, Fimland MS, Mosti MP, Wang E. Strength training-induced responses in older adults: Attenuation of descending neural drive with age. *Age (Dordr)* 37: 9784, 2015.
620. Unhjem R, Nygard M, van den Hoven LT, et al. Lifelong strength training mitigates the age-related decline in efferent drive. *J Appl Physiol (1985)* 121: 415–423, 2016.
621. Valenzuela T. Efficacy of progressive resistance training interventions in older adults in nursing homes: A systematic review. *J Am Med Directors Assoc* 13: 418–428, 2012.
622. Valkeinen H, Hakkinen K, Pakarinen A, et al. Muscle hypertrophy, strength development, and serum hormones during strength training in elderly women with fibromyalgia. *Scand J Rheumatol* 34: 309–314, 2005.
623. Vaughan S, Wallis M, Polit D, Steele M, Shum D, Morris N. The effects of multimodal exercise on cognitive and physical functioning and brain-derived neurotrophic factor in older women: A randomised controlled trial. *Age Ageing* 43: 623–629, 2014.
624. Vellera HL, Kleeberger SR, Lightfoot JT. Inter-individual variation in adaptations to endurance and resistance exercise training: Genetic approaches towards understanding a complex phenotype. *Mamm Genome* 29: 48–62, 2018.
625. Venturelli M, Lanza M, Muti E, Schena F. Positive effects of physical training in activity of daily living-dependent older adults. *Exp Aging Res* 36: 190–205, 2010.
626. Vettor R, Milan G, Franzin C, et al. The origin of intermuscular adipose tissue and its pathological implications. *Am J Physiol Endocrinol Metab* 297: E987–E998, 2009.
627. Villareal DT, Smith GL, Sinacore DR, Shah K, Mittendorfer B. Regular multicomponent exercise increases physical fitness and muscle protein anabolism in frail, obese, older adults. *Obesity (Silver Spring)* 19: 312–318, 2011.
628. Villareal DT, Steger-May K, Schechtman KB, et al. Effects of exercise training on bone mineral density in frail older women and men: A randomised controlled trial. *Age Ageing* 33: 309–312, 2004.

629. Vincent KR, Braith RW, Feldman RA, et al. Resistance exercise and physical performance in adults aged 60 to 83. *J Am Geriatr Soc* 50: 1100–1107, 2002.
630. Vincent KR, Vincent HK, Braith RW, Lennon SL, Lowenthal DT. Resistance exercise training attenuates exercise-induced lipid peroxidation in the elderly. *Eur J Appl Physiol* 87: 416–423, 2002.
631. Vingren JL, Kraemer WJ, Ratamess NA, et al. Testosterone physiology in resistance exercise and training: The up-stream regulatory elements. *Sports Med* 40: 1037–1053, 2010.
632. Visser M, Goodpaster BH, Kritchevsky SB, et al. Muscle mass, muscle strength, and muscle fat infiltration as predictors of incident mobility limitations in well-functioning older persons. *J Gerontol A Biol Sci Med Sci* 60: 324–333, 2005.
633. Visser M, Pahor M, Taaffe DR, et al. Relationship of interleukin-6 and tumor necrosis factor- α with muscle mass and muscle strength in elderly men and women: The health ABC study. *J Gerontol A Biol Sci Med Sci* 57: M326–M332, 2002.
634. Visser M, Simonsick EM, Colbert LH, et al. Type and intensity of activity and risk of mobility limitation: The mediating role of muscle parameters. *J Am Geriatr Soc* 53: 762–770, 2005.
635. Volkert J, Schulz H, Harter M, Włodarczyk O, Andreas S. The prevalence of mental disorders in older people in Western countries—A meta-analysis. *Ageing Res Rev* 12: 339–353, 2013.
636. Walker S, Santolamazza F, Kraemer W, Hakkinen K. Effects of prolonged hypertrophic resistance training on acute endocrine responses in young and older men. *J Aging Phys Act* 23: 230–236, 2015.
637. Walston J, Fried LP. Frailty and the older man. *Med Clin North Am* 83: 1173–1194, 1999.
638. Webber SC, Porter MM, Menec VH. Mobility in older adults: A comprehensive framework. *Gerontologist* 50: 443–450, 2010.
639. Weiss CO. Frailty and chronic diseases in older adults. *Clin Geriatr Med* 27: 39–52, 2011.
640. Welle S, Thornton C, Jozefowicz R, Statt M. Myofibrillar protein synthesis in young and old men. *Am J Physiol* 264: E693–E698, 1993.
641. Welle S, Thornton C, Statt M. Myofibrillar protein synthesis in young and old human subjects after three months of resistance training. *Am J Physiol* 268: E422–E427, 1995.
642. West S, King V, Carey TS, et al. Systems to rate the strength of scientific evidence. *Evid Rep Technol Assess (Summ)* 47: I–II, 2002.
643. Westcott WL. Resistance training is medicine: Effects of strength training on health. *Curr Sports Med Rep* 11: 209–216, 2012.
644. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: A report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *J Am Coll Cardiol* 71: e127–e248, 2018.
645. Whitney DG, Singh H, Miller F, et al. Cortical bone deficit and fat infiltration of bone marrow and skeletal muscle in ambulatory children with mild spastic cerebral palsy. *Bone* 94: 90–97, 2017.
646. WHO. International Classification of Functioning, Disability, and Health: ICF. Geneva, Switzerland: World Health Organization, 2001.
647. Williams MA, Haskell WL, Ades PA, et al. Resistance exercise in individuals with and without cardiovascular disease: 2007 update: A scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, physical activity, and metabolism. *Circulation* 116: 572–584, 2007.
648. Williams MA, Stewart KJ. Impact of strength and resistance training on cardiovascular disease risk factors and outcomes in older adults. *Clin Geriatr Med* 25: 703–714, 2009; ix.
649. Wong E, Backholer K, Gearon E, et al. Diabetes and risk of physical disability in adults: A systematic review and meta-analysis. *Lancet Diabetes Endocrinol* 1: 106–114, 2013.
650. Wood RH, Reyes R, Welsch MA, et al. Concurrent cardiovascular and resistance training in healthy older adults. *Med Sci Sports Exerc* 33: 1751–1758, 2001.
651. Wright NC, Looker AC, Saag KG, et al. The recent prevalence of osteoporosis and low bone mass in the United States based on bone mineral density at the femoral neck or lumbar spine. *J Bone Miner Res* 29: 2520–2526, 2014.
652. Writing Group M, Mozaffarian D, Benjamin EJ, et al. Heart disease and stroke statistics-2016 update: A report from the American heart association. *Circulation* 133: e38–360, 2016.
653. Wu Y, Wang W, Liu T, Zhang D. Association of grip strength with risk of all-cause mortality, cardiovascular diseases, and cancer in community-dwelling populations: A meta-analysis of prospective cohort studies. *J Am Med Dir Assoc* 18: 551.e17–551.e35, 2017.
654. Xue QL. The frailty syndrome: Definition and natural history. *Clin Geriatr Med* 27: 1–15, 2011.
655. Xue QL, Beamer BA, Chaves PH, Guralnik JM, Fried LP. Heterogeneity in rate of decline in grip, hip, and knee strength and the risk of all-cause mortality: The women's health and aging study II. *J Am Geriatr Soc* 58: 2076–2084, 2010.
656. Yamada M, Arai H, Sonoda T, Aoyama T. Community-based exercise program is cost-effective by preventing care and disability in Japanese frail older adults. *J Am Med Dir Assoc* 13: 507–511, 2012.
657. Yang Z, Scott CA, Mao C, Tang J, Farmer AJ. Resistance exercise versus aerobic exercise for type 2 diabetes: A systematic review and meta-analysis. *Sports Med* 44: 487–499, 2014.
658. Yarasheski KE, Zachwieja JJ, Campbell JA, Bier DM. Effect of growth hormone and resistance exercise on muscle growth and strength in older men. *Am J Physiol* 268: E268–E276, 1995.
659. Yue GH, Ranganathan VK, Siemionow V, Liu JZ, Sahgal V. Older adults exhibit a reduced ability to fully activate their biceps brachii muscle. *J Gerontol A Biol Sci Med Sci* 54: M249–M253, 1999.
660. Zanusso S, Sieverdes JC, Smith N, Carraro A, Bergamin M. The effect of a strength training program on affect, mood, anxiety, and strength performance in older individuals. *Int J Sport Psychol* 43: 53–66, 2012.
661. Zhang P, Peterson M, Su GL, Wang SC. Visceral adiposity is negatively associated with bone density and muscle attenuation. *Am J Clin Nutr* 101: 337–343, 2015.
662. Zoico E, Rossi A, Di Francesco V, et al. Adipose tissue infiltration in skeletal muscle of healthy elderly men: Relationships with body composition, insulin resistance, and inflammation at the systemic and tissue level. *J Gerontol A Biol Sci Med Sci* 65: 295–299, 2010.
663. Zwijssen SA, Depla MF, Niemeijer AR, Francke AL, Hertogh CM. The concept of restraint in nursing home practice: A mixed-method study in nursing homes for people with dementia. *Int Psychogeriatr* 23: 826–834, 2011.

